

Supplementary Table 2. Depression-Associated SNP Identified in the WLS

Gene	Encodes	SNP	Alleles	Chr#/Location	Residue	Associated disease/behavior
ACVR2B	activin receptor IIB	rs3749386	T/C	3/intron 1	--	left-right axis malformations*(1)
APOC3	apolipoprotein C-III	rs2854116	T/C	11/promoter (-455)	--	nonalcoholic fatty liver disease(2)
<i>DRD2/ANKK1</i>	dopamine receptor D2/ankyrin repeat and kinase domain containing 1	rs1800497	C/T	11/exon (ANKK1)	Glu713Lys	obesity, drug addiction (3)
<i>DRD2</i>	dopamine receptor D2	rs2242592	T/C	11/3'	--	schizophrenia (4)
<i>FTO</i>	fat mass and obesity associated	rs1421085	T/C	16/intron 1	--	obesity (5-7), mental disorders (8)
<i>GNRH1</i>	gonadotropin-releasing hormone	novel SNP	T/C	8/promoter	--	Alzheimer's disease (9)
<i>IL6</i>	interleukin 6	rs1800795	C/G	7/promoter (-174)	--	arthritis (10), breast cancer (11), diabetes (12), depression (13)

Gene association only

References

1. Kosaki R, Gebbia M, Kosaki K, Lewin M, Bowers P, Towbin JA, et al. Left-right axis malformations associated with mutations in ACVR2B, the gene for human activin receptor type IIB. *American Journal of Medical Genetics.* 1999;82(1):70-6.
2. Petersen KF, Dufour S, Hariri A, Nelson-Williams C, Foo JN, Zhang X-M, et al. Apolipoprotein C3 Gene Variants in Nonalcoholic Fatty Liver Disease. *New England Journal of Medicine.* 2010;362(12):1082-9.
3. Blum K, Braverman ER, Wood RC, Gill J, Li C, Chen TJ, et al. Increased prevalence of the Taq I A1 allele of the dopamine receptor gene (DRD2) in obesity with comorbid substance use disorder: a preliminary report. *Pharmacogenetics.* 1996 Aug;6(4):297-305.
4. Dubertret C, Bardel C, Ramoz N, Martin P-M, Deybach J-C, Adès J, et al. A genetic schizophrenia-susceptibility region located between the ANKK1 and DRD2 genes. *Progress in Neuro-Psychopharmacology and Biological Psychiatry.* 2010;34(3):492-9.
5. Dina C, Meyre D, Gallina S, Durand E, Korner A, Jacobson P, et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet.* 2007;39(6):724-6.
6. Tung Y-CL, Yeo GSH. From GWAS to biology: lessons from FTO. *Annals of the New York Academy of Sciences.* 2011;1220(1):162-71.
7. Chabris C, Hebert B, Benjamin D, Beauchamp J, Cesari D, van der Loos M, et al. Most reported genetic associations with general intelligence are probably false positives. *Psychological Science.* 2011;In press.
8. Kivimaki M, Jokela M, Hamer M, Geddes J, Ebmeier K, Kumari M, et al. Examining Overweight and Obesity as Risk Factors for Common Mental Disorders Using Fat Mass and Obesity-Associated (FTO) Genotype-Instrumented Analysis. *Am J Epidemiol.* 2011 Feb 15;173(4):421-9.
9. Wilson AC. Investigations into gonadotropin-releasing hormone receptor signaling in extrahypothalamic neurons of the brain: Implications for normal cognition and the pathogenesis of Alzheimer's disease: Thesis. University of Wisconsin, Madison, USA; 2009.
10. Fishman D, Faulds G, Jeffery R, Mohamed-Ali V, Yudkin JS, Humphries S, et al. The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *The Journal of Clinical Investigation.* 1998;102(7):1369-76.
11. Slattery ML, Curtin K, Sweeney C, Wolff RK, Baumgartner RN, Baumgartner KB, et al. Modifying Effects of IL-6 Polymorphisms on Body Size-Associated Breast Cancer Risk. *Obesity.* 2008;16(2):339-47.
12. Huth C, Heid IM, Vollmert C, Gieger C, Grallert H, Wolford JK, et al. IL6 Gene Promoter Polymorphisms and Type 2 Diabetes. *Diabetes.* 2006 October 2006;55(10):2915-21.
13. Bull SJ, Huezo-Diaz P, Binder EB, Cubells JF, Ranjith G, Maddock C, et al. Functional polymorphisms in the interleukin-6 and serotonin transporter genes, and depression and fatigue induced by interferon-alpha and ribavirin treatment. *Mol Psychiatry.* 2009 Dec;14(12):1095-104.