

Table S3. Toxicological parameters

	PBOISL				PBOISNC			
	AST (mg/dL)	ALT (mg/dL)	BUN (mg/dL)	CRE (mg/dL)	AST (mg/dL)	ALT (mg/dL)	BUN (mg/dL)	CRE (mg/dL)
Saline	60 ± 23.6	21 ± 4.2	36.0 ± 5.3	1.1 ± 0.05	89 ± 37.6	45 ± 18.9	41.4 ± 5.4	1.1 ± 0.07
1mg/kg	60 ± 24.8	22 ± 9.7	38.0 ± 4.9	1.1 ± 0.02	60 ± 22.7	23 ± 9.5	41.8 ± 10.0	1.1 ± 0.12
5mg/kg	69 ± 58.4	19 ± 3.8	45.0 ± 8.0	1.0 ± 0.09	65 ± 26.2	36 ± 16.2	38.8 ± 4.5	1.0 ± 0.05
10mg/kg	43 ± 7.2	16 ± 3.8	50.7 ± 10.1 ^a	1.0 ± 0.01	58 ± 12.6	26 ± 8.1	43.9 ± 9.2	1.1 ± 0.05
20mg/kg	55 ± 7.7	34 ± 14.6	48.1 ± 16.0	1.0 ± 0.05				

Blood samples were collected from mice tail veins after 6 weeks of administration.

^aValues in the line with superscripts indicate significant differences ($P < 0.05$) (\pm SD).