

Table 3: Unadjusted temperature effects on ambulance attendances in Brisbane, Australia. Effects given as percent change per 1 °C above or below threshold (Threshold stated in brackets for each category)

	Lag in days	Attendance category (95% C.I)			
		Total (22 °C)	Cardiovascular (22 °C)	Respiratory (22 °C)	Other (15.5/28 °C)
Heat effect	0-1	1.51 (0.54, 2.49)	0.81 (-0.15, 1.77)	0.29 (-0.93, 1.51)	23.52 (13.23, 34.31)
	2-15	-0.49 (-1.96, 0.99)	-1.25 (-2.70, 0.21)	-1.16 (-3.00, 0.69)	-27.55 (-53.20, -1.41)
	16-27	1.02 (-0.21, 2.24)	1.12 (-0.09, 2.33)	1.89 (0.36, 3.43)	-0.95 (-23.62, 22.16)
Cold effect	0-1	-0.57 (-1.27, 0.14)	-0.68 (-1.36, 0.01)	-0.03 (-0.81, 0.75)	-3.52 (-6.61, -0.39)
	2-15	0.86 (-0.22, 1.93)	1.71 (0.66, 2.76)	3.83 (2.63, 5.02)	-0.97 (-5.82, 3.89)
	16-27	1.10 (0.26, 1.93)	0.08 (-0.73, 0.90)	2.79 (1.86, 3.72)	5.81 (1.60, 10.03)