

The nucleotide sequence of 3-isopropylmalate dehydrogenase gene from *Bacillus caldovenax*T.Sekiguchi, M.Suda, T.Ishii, Y.Nosoh and K.Tsuda¹

Department of Life Science, Faculty of Science, Tokyo Institute of Technology, Nagatsuta, Yokohama, Kanagawa 227 and ¹Institute of Polymers and Textiles, Yatabe-Higashi, Tsukuba, Ibaraki 305, Japan

Submitted December 30, 1986

FEATURES: The gene coding for 3-isopropylmalate dehydrogenase from *Bacillus coagulans* (facultative thermophile) was sequenced (1). Since the enzyme gene from *B. caldovenax* (obligate, moderate thermophile) had been cloned in *Escherichia coli* using pBR322 as a vector (2), the nucleotide sequence for the *B. caldovenax* enzyme gene was determined.

```

ATGGGAAACTCGGATTGCCCTTCGCCAGGGGATGCCATGCCAAAAGAAGTGACGCCGAGGGTCCGAGGTTAAAAAGCGTCGCCATCCGGTTGGCACCAGTTACGTTGAA 120
M G N Y R I A V L P G D G I G K E V T S G A V E V L K A V G I R F G H E F T F E
TATGGATTGATCGGTGAGCGGGCATCGATGAAGCCGGGAGCCGGCTCCCGAGAAAACCGTCCGCCTTCCGAGAAAGCGATCCGGTCTGCTTGGGGCGGTCGGCGGAGTGG 240
Y G L I G G A A I D E A G T P L P E E T V R L C R E S D A V L L G A V G G P K W
GATGACACCCCTCCGATTCCGCCGAAAGGATTGCTTCCCAATCGCAAACGACTCCGACTTATGCCAACCTGGGCCCGGTGTTTGCTATGACAGTTGGTTCCCGTTCGGC 360
D D N F P H L R P E K G L L A I R K O L D L Y A N L R P V V C Y D S L V S R S P
TTAAACACCTCTCGTCAAGGGCTCGATTTCATCTCCCGAACCTGACCCCGCCATTACTTCCCAAACCCACCCCGTCCTCGAAAACGGCGAGAAAAACGCACTGGACACC 480
L K P D L V Q G V D F V I V R E L T G G I Y F G Q P S A V V E N G E E K A V D T
TTGTTATATAAAAGGAGAAATTAAGCGGATCGTCCCGATGGCTTGTGAGCTCGCACCGGGCGAGGAAAAAAAGTGAATTCGGTCGACAAGCCGAACGTCCTTTCGTCAGCCCTCA 600
L L Y K K E E I E R V R M A F E L A R G R R K K V T S V D K A H V L S S S R L
TGGAGGGAAAGTGGCCGAAGAGGTGGVCCAGCAGGAGTTCGGGACGTCACCGTGGACCATATGCTTGTGACATGGGGATGCAAGCTCATCCGTGGCCGAACCCAGTTGAC 720
W R E V A E E V A N E F P D V T L E H M L V D M R M O L I R A P K O F D V I V T
GAAAAACATTTGGCGACATTTAAAGCGATGAAGCTCCATGTTGTCGGGTTGCTGGGATGCTGGCGTCCGCCAGCTTATGCCCTTCCGTCGAGCTTGTATGACCCGTGACCGT 840
E H M F G D I L S D E A S M L S G S L G M L P S A S L S A G S P S L Y E P V H G
TCAGCGGCTGATATGCCCGCATGAAACAAACCGCAATCCGGATTGGCCGATGTCGGCCGATGACTTCGACTGCTGGCTGGGATGACCCGAACCCAGTTGACCCGTGAC 960
S A P D I A G M N K A A M P I A A I L S A A M H L R L S F G L T A E A G G R A R V
TGGCAGGGCTGCCCTAGGTCTGGCAGGGACTTGGCCAACGGCGGGCCATCTATCCACGAAGAAATGGCGAAGAAATCAAAGCGGGGCTGGCTGGATTATAACGCCATTGCTAA 1080
W O A L A L G S G S R L G O R R P H L S T R E M V E E I K A A V L D Y T A I A O
ATTATGACCGTTATGCTAA 1101
I M T V Y A *

```

COMMENTS: The G+C contents of coding region and third letter of the codon for the *B. caldovenax* enzyme were 56.7 and 65.2%, respectively. The values were intermediate between those for the *B. coagulans* (53.1 and 55.7%, respectively (1) and for the *Thermus thermophilus* enzyme (70.0 and 89.4%, respectively).

REFERENCES:

1. Sekiguchi, T., et al., (1986) Biochim. Biophys. Acta 867, 36-44.
2. Sekiguchi, T., et al., (1986) FEMS Microbiol. Lett. 36, 41-45.
3. Kagawa, Y., et al., (1984) J. Biol. Chem. 259, 2956- 960.