

**Table S1 Quantitation of Proteins Shown in Figure 5D.**

		C	T	HcyL	RL1	RL2	HRL	E1	E2	ERL
nSREBP1c	Average	0.1130	0.1665	0.0376	0.3636	0.2480	0.0441	0.1869	0.1673	0.4488
	SD	0.0146	0.0094	0.0171	0.1245	0.0690	0.0181	0.0448	0.0591	0.0722
	p value 1		0.0059	0.0044	0.0258	0.0295	0.0068	0.0531	0.1974	0.0014
	p value 2									0.0059
PGC1	Average	0.5778	0.2880	0.0389	0.4618	0.4572	0.0825	0.3516	0.3788	0.1212
	SD	0.0765	0.0450	0.0225	0.0689	0.1574	0.0474	0.0658	0.1080	0.0510
	p value 1		0.0048	0.0003	0.1226	0.2984	0.0007	0.0178	0.0598	0.0010
	p value 2									0.0087
Derlin	Average	0.6131	0.4378	0.2504	0.6319	0.6859	0.1442	0.5031	0.4725	0.4346
	SD	0.0179	0.0902	0.0736	0.0683	0.1129	0.1101	0.0530	0.1298	0.0943
	p value 1		0.0298	0.0012	0.6700	0.3323	0.0019	0.0271	0.1365	0.0322
	p value 2									0.3342
CYP2E1	Average	1.4986	1.7819	1.2681	0.4517	0.6180	2.0467	2.5888	1.8917	1.1952
	SD	0.1429	0.2309	0.2084	0.2204	0.2731	0.5343	0.5237	0.4534	0.2746
	p value 1		0.1451	0.5208	0.0023	0.0078	0.1612	0.0254	0.2254	0.1648
	p value 2									0.0151
SERCA	Average	2.9375	2.0278	1.3887	2.0116	2.1028	1.3627	2.5337	2.7768	1.8318
	SD	0.2340	0.3871	0.2275	0.3305	0.4843	0.2489	0.2018	0.5258	0.3128
	p value 1		0.0253	0.0012	0.0167	0.0548	0.0013	0.0863	0.6539	0.0080
	p value 2									0.5315

C, control; T, tunicamycin; HcyL, homocysteine thiolactone; RL, ritonavir and lopinavir at 10 µg/ml (RL1) and 5 µg/ml (RL2); HRL, homocysteine thiolactone plus RL at 5 µg/ml; E1, alcohol (85 mM); E2, alcohol (35 mM); ERL, RIT and LOP (10 µg/ml) plus alcohol at 85 mM; Protein quantity was normalized with Cyclin D; Comparisons were made between control and individual treatments (p value 1) or between RL1 and ERL (p value 2).