

ESM Table 1

Table 1. CAC and SMPC levels *in vitro* expressed as 10^3 cells per 10^6 WBC and per mL blood.

		Non-Type 2 diabetes					Type 2 diabetes				
		all	no MVD	MVD	PAD	CAD	all	no MVD	MVD	PAD	CAD
n		24	7	17	10	7	26	7	19	10	9
CAC	x 10^3 per 10^6 WBC	10.5 ± 1.4	11.3 ± 2.0	10.2 ± 1.8	13.3 ± 2.6	5.7 ± 1.5	8.3 ± 0.9	11.2 ± 1.9 ^a	7.0 ± 0.9	5.9 ± 0.9	8.6 ± 1.6
	x 10^3 per mL blood	77.8 ± 14.1	60.3 ± 11.9	84.9 ± 19.3	120.0 ± 27.8 ^b	35.0 ± 7.2	61.0 ± 6.2	78.6 ± 13.2	54.6 ± 6.6	49.7 ± 9.2	55.7 ± 10.6
SMPC	x 10^3 per 10^6 WBC	33.3 ± 4.2	17.2 ± 3.3 ^c	39.9 ± 5.0	44.7 ± 6.9	33.0 ± 6.8	29.9 ± 2.7	29.3 ± 4.9	30.2 ± 3.3	32.2 ± 4.9	28.0 ± 4.6
	x 10^3 per mL blood	250.0 ± 44.1	94.1 ± 20.8 ^d	318.8 ± 57.2	407.8 ± 83.4	191.6 ± 41.5	232.2 ± 32.0	235.3 ± 85.7	231.2 ± 33.9	275.8 ± 56.7	181.7 ± 29.7

Data presented as mean ± SEM.

Statistically significant with ANOVA compared to:

^aT2DM no MVD compared to T2DM with MVD (p<0.05)

^bNon-T2DM with PAD compared to non-T2DM with CAD (p<0.05)

^cHealthy compared with T2DM, non-T2DM with MVD and non-T2DM with PAD (p<0.05).

^dHealthy compared with T2DM (p<0.05), non-T2DM with MVD (p<0.05) and non-T2DM with PAD (p<0.01)