

Supplementary Table 4. Distribution of STs with significant admixture among BAPS groups and source of isolation

BAPS groups	BAPS subgroups	Number (%) of STs identified in the different source groups with significant admixture												No (%) of STs with admixture in
		Hosp patients		Non-hosp persons		Pets		Pigs		Poultry		Calves		
		Admixture	No admixture	Admixture	No admixture	Admixture	No admixture	Admixture	No admixture	Admixture	No admixture	Admixture	No admixture	
1		1 (4%)	24	0	9	0	0	1 (10%)	9	0	0	0	0	2 (4%)
2	1	2 (3%)	64	0	11	0	12	1 (5%)	18	0	44	0	0	3 (2%)
	2	1 (100%)	0	0	0	0	0	0	0	0	0	0	0	1 (100%)
	3	0	32	1 (12%)	7	0	7	0	6	0	3	0	0	1 (2%)
	4	4 (100%)	0	0	0	1 (100%)	0	1 (100%)	0	0	3	0	0	8 (53%)
	Total	7 (7%)	96	1 (5%)	18	1 (5%)	19	1 (4%)	24	0	50	0	0	13 (6%)
3	1	0	31	0	5	0	4	0	6	0	1	0	6	0 (0%)
	2	6 (100%)	0	3 (100%)	0	3 (100%)	0	7 (100%)	0	1 (100%)	0	0	0	17 (100%)
	3	2 (3%)	71	0	6	1 (17%)	5	0	6	0	9	0	3	3 (3%)
	4	1 (100%)	0	3 (100%)	0	0	0	1 (100%)	0	0	0	0	0	4 (100%)
	5	4 (31%)	9	0	1	0	3	0	0	2 (33%)	2	0	0	5 (24%)
	Total	13 (10%)	111	6 (30%)	12	4 (25%)	12	8 (40%)	12	2 (14%)	12	0	9	29 (15%)
4		1	5	0	1	0	0	0	0	0	0	0	0	2 (22%)
5		2	3	0	0	0	0	1 (50%)	1	1 (50%)	1	0	0	3 (33%)
6		0	2	0	0	0	0	0	0	0	0	0	0	0 (0%)
7		0	7	0	0	0	0	0	0	0	0	0	0	0 (0%)
Total		24 (9%)	248	7 (15%)	40	5 (14%)	31	12 (21%)	46	3 (5%)	63	0	9	