

**Table S1. Sequence of the oligonucleotides used for the construction of pJLE-derived plasmids**

<b>1</b>	5' GATCT <b>G</b> GGCCATTACGGCCTCTAGATTAGGCCGCCTCGGCCGGTAC 3'
<b>2</b>	5' <u>CGGCCGAGGCGGCCT</u> AATCTAGAGGCCGTAATGGCCA 3'
<b>3</b>	5' GATCT <b>G</b> GGCCATTACGGCCTCTAGATAAGGCCGCCTCGGCCGGTAC 3'
<b>4</b>	5' <u>CGGCCGAGGCGGCCT</u> TATCTAGAGGCCGTAATGGCC <b>G</b> CA 3'
<b>5</b>	5' GATCT <u>GGCCATTACGGCCT</u> CTAGATAAGGCCGCCTCGGCCGGTAC 3'
<b>6</b>	5' <u>CGGCCGAGGCGGCCT</u> TATCTAGAGGCCGTAATGGCCA 3'
<b>7</b>	5'GATCTGGTGGCGGTTCTGGCGGTGGTTCTGGTGGCGGTTCCGGCGGTGGTT CTGGCGGTGGCTCTGGTGGCGGCTCCGGTGGTGGTAGCTCTGGCGGTA 3'
<b>8</b>	5'GATCTACCGCCAGAGCTACCACCACCGGAGCCGCCACCAGAGCCACCGCC AGAACCACCGCCGGAACCGCCACCAGAACCACCGCCAGAACCAGCCACCA 3'

Underlined nucleotides are *Sfi*I restriction sites; bold nucleotides are additional nucleotides that allow expression of the cDNA in distinct reading-frames.