

Supplemental Material

Modelling phenoxy acid herbicide mineralization and growth of microbial degraders in 15 soils monitored by quantitative real-time PCR of the functional *tfdA* gene

Jacob Bælum^{1,2}, Emmanuel Prestat^{3,4}, Maude M. David³, Bjarne W. Strobel⁵, and Carsten S. Jacobsen^{1,5,6*}

¹Geological Survey of Denmark and Greenland (GEUS), Department of Geochemistry, Øster Voldgade 10, DK-1350 Copenhagen K, Denmark; ²Technical University of Denmark, Novo Nordisk Foundation Center for Biosustainability, 2970 Hørsholm, Denmark; ³Lawrence Berkeley National Laboratory, Earth Sciences Division, Berkeley, California 94720, USA; ⁴Kansas State University, Division of Biology, Manhattan, Kansas 66506, USA; ⁵University of Copenhagen, Department of Basic Sciences and Environment, Thorvaldsensvej 40, DK-1871 Frederiksberg C, Denmark. ; ⁶Center for Permafrost , CENPERM, University of Copenhagen, Øster Voldgade 10, DK-1350 Copenhagen K, Denmark

Contains 2 figures.




Figure S1: Mineralization curves for all of the sixteen soil scenarios. The mineralization curves of 2,4-D, MCPA, and MCPP are given for the individual soils (panel A-P). Error bars represent standard errors of the mean for soil triplicates




Figure S2. Quantification of *rpoB* genes. Error bars represent standard errors of the mean for soil triplicates.