

1
2
3
4
5

Supplementary TABLE 2. Polymorphisms found on the full-length nucleotide sequence of the *cqm1*_{REC-D16} allele from the *Culex quinquefasciatus* *cqm1* gene (GenBank accession number DQ333335). Diverging nucleotides which cause amino acids changes in deduced sequence coded by *cqm1* (Romão et al., FEBS J. 273:1556-1568) are marked (*).

Nucleotide position	<i>cqm1</i>	<i>cqm1</i> _{REC-D16}	Nucleotide position	<i>cqm1</i>	<i>cqm1</i> _{REC-D16}
7	C	A*	736	T	C*
15	A	T	1084	G	A
22	A	G*	1140	C	T
25	C	T	1262	C	T
44	T	C*	1271	A	G
120	T	C	1325	T	C
123	G	A	1328	C	A
156	T	A	1340	C	A
249	G	C	1349	A	C
252	A	G	1367	T	C
264	T	C	1391	G	A
284	C	A*	1439	T	G
288	C	T	1505	A	T
321	C	T	1506	T	C
337	T	G*	1559	C	T
357	G	A	1577	G	A
369	T	C	1619	C	T

393	C	T	1749	C	T
422	C	T	1772	C	T
438	C	T	1776	T	A
462	C	T	1839	T	C
611	C	T*	—	—	—

6

7