

Microsatellite markers for the characterization of native and introduced populations of *Plasmopara viticola*, the causal agent of grapevine downy mildew

Mélanie Rouxel, Daciana Papura, Marilise Nogueira, Virginie Machefer, Damien Dezette, Sylvie Richard-Cervera, Sébastien Carrere, Pere Mestre, and François Delmotte

Table S1: Date of collection and geographical origin of *Plasmopara viticola* strains and populations analysed in this study.

Sample name	Number of isolates	Year	Location	Country	GPS position	
Strains						
Pv221	1	2009	Blanquefort	France	N 44° 55' 14.4336"	W 0° 38' 30.5916"
SC	1	2005	Colmar	France	N 48° 3' 51.3396"	E 7° 20' 2.5044"
SL	1	2008	Lednice	Czech Republic	N 48° 47' 28.6044"	E 16° 47' 42.4278"
Populations						
Gironde	16	2003	Sauternes	France	N 44° 31' 59.466"	W 0° 19' 58.1874"
Rhine Valley	16	2003	Freibourg	Germany	N 47° 58' 27.6672"	E 7° 48' 16.704"
Vaucluse	16	2005	Sorgues	France	N 44° 1' 51.0846"	E 4° 51' 41.8968"
Michigan	16	2010	Fennville	USA	N 42° 35' 59.9208"	W 86° 9' 13.7514"
Virginia	16	2010	Hidden Brook	USA	N 39° 12' 18.6366"	W 77° 29' 57.861"
New York	16	2010	Geneva	USA	N 42° 52' 12.525"	W 77° 2' 38.5368"