

## Supplementary Data

**SUPPLEMENTARY TABLE S1. DEMOGRAPHICS, CLINICAL CHARACTERISTICS, AND HARDY–WEINBERG EQUILIBRIUM OF THE GENOTYPES OF *CYP27B1*-1260 AND *VDR* TAQ I IN HEPATITIS B VIRUS PATIENTS, HEPATITIS B VIRUS INFECTION RESOLVERS, AND HEALTHY CONTROLS**

	<i>Patients</i> (n=274)	<i>Resolvers</i> (n=68)	<i>Controls</i> (n=158)	p
Sex (male/female)	210/64	43/25	118/40	0.076
Age (year, mean±SD)	45.51±13.28	46.69±13.72	43.11±13.28	0.099
HWE (observed vs. expected)				
<i>CYP27B1</i> -1260				
<i>p</i> -value	0.984	0.537	0.644	
<i>VDR</i> Taq I				
<i>p</i> -value	0.485	1.000	0.985	
HBeAg status		—	—	
HBeAg(+)	112			
HBeAg(−)	162			
Clinical diagnosis		—	—	
Chronic hepatitis	94			
Liver cirrhosis	90			
HCC	90			

HWE, Hardy–Weinberg equilibrium; HCC, hepatocellular carcinoma.