

Members of the germin-like protein family in *Brassica napus* are candidates for the initiation of an oxidative burst that impedes pathogenesis of *Sclerotinia sclerotiorum*.

Steffen Rietz, Friederike Bernsdorff and Daguang Cai

Table S1. Primer combinations for qPCR analysis of the *BnGLP* gene family and tubulin as reference gene.

	Forward 5'-3'	Reverse 5'-3'
<i>BnGLP1</i>	TGATGGGTACCTCCATGTCTCGT	GGCAGGAGCATTGTTAGGGTTCT
<i>BnGLP2</i>	CCGTCAATGTTAGCAACCTTCCA	GTTTTGGTGAAGAGGCGATTCC
<i>BnGLP3</i>	GGTTTCTCACCACCGCCAATAG	AACCGAACAGAGTAGCGCCAAG
<i>BnGLP4</i>	ATTTCTTCTTTGCGGGCATCAG	CTCGCCTTCCAACACGAAGATT
<i>BnGLP5</i>	AATTCTGTAAGGATCCGAAGCAAGC	TCGATTAGGACAAGGATCTCAGTGC
<i>BnGLP6</i>	TGGTCCTGGCGACATGTTTG	GGAATCGCTGGTTTCGCAAG
<i>BnGLP7</i>	GCTGATCTCAAACCCGGAGCTA	CCGGGGCTTTGGCTATTAAGTC
<i>BnGLP8</i>	ACCCTGACCAAGTCACCGAAAAT	TGTGAAGAGGAATGACACCACCA
<i>BnGLP9</i>	GTAGCTGATCCGAGCGGTCTT	CCAGCGGCTAAGTCAAGACGA
<i>BnGLP10</i>	CAAGACTTCTGTGTGGCCAACCT	AGTGTGCATTGGGATCACACCTT
<i>BnGLP11</i>	CTCCTCCTCATTACCTCACCAA	ATTGTGTTGAGACCCGGGAGATT
<i>BnGLP12</i>	TCCCTCTCATCTGACCCTGACC	GGAGCGAACTCAACGTTGTTCA
<i>BnGLP13</i>	ACGTCGGGTTTGTCTTCCAA	ACACAGCGTTTTCGATTGTGAT
<i>BnGLP14</i>	TTCGTAACAACCGCAGGGAAA	GCGACTGTGTTCCAGGCAACT
<i>Tubulin</i>	AGGTCTCCGACACTGTTGTTG	GGAGTTGAGTTGACCAGGGA

Members of the germin-like protein family in *Brassica napus* are candidates for the initiation of an oxidative burst that impedes pathogenesis of *Sclerotinia sclerotiorum*.

Steffen Rietz, Friederike Bernsdorff and Daguang Cai

Table S2. Primer combinations for full length gene cloning into the pDONR201 Gateway vector.

		5' - 3'
<i>BnGLP3</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGGCTTCACCAACTTCATT
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTCTTTCTTGGGCTGGAACCTGC
<i>BnGLP7</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGTTTCTTACAAAAGCTCTTC
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTTCATGTTTCATTAGACATGACAG
<i>BnGLP8</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGAAGATGATAATTCACAT
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTTCATCAGTTCCTCCAAGCACAC
<i>BnGLP10</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGTTGCGCACTATCTTCCT
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTTCATGAGTCCCAAGTATCTTTT
<i>BnGLP12</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGGCTTCTCTTTTCACATC
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTCAACAGCAAACCTTGGACTTGAG
<i>gf2.8</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGGGGTACTCCAAAACCCTAGT
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTCAAACCCAGCGGCAAACCTTGGACT
<i>GFP</i>	forward	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGGTGAGCAAGGGCGAGGA
	reverse	GGGGACCACTTTGTACAAGAAAGCTGGGTCTTGTACAGCTCGTCCATGC

Figure S1




Figure S1. Differential transcription of the β -tubulin reference gene between mock and *S. sclerotiorum* treated leaf samples of *B. napus* 'Falcon' and 'Zhongshuang 9'. Quantitative RT-PCR was used to determine cycle threshold (C_t) values of β -tubulin as a reference to normalize *BnGLP* gene transcript level in Fig. 2B and C. Shown are the differences in β -tubulin C_t values between 6 h *S. sclerotiorum* and 6 h mock treatments. Columns represent the averages of four independent experiments with bars depicting corresponding standard errors.