

Table S2 Plasmids used in this study.

Plasmid	Genotype	Source
dRed-HDEL	<i>trp1::DsRED-HDEL:TRP1</i> integration plasmid	(BEVIS <i>et al.</i> 2002)
pBS35	<i>mCHERRY/HYGB</i> integration plasmid	(SHANER <i>et al.</i> 2004)
pRS315	CEN/LEU2	(SIKORSKI and HIETER 1989)
pRS425	2μ/LEU2	(CHRISTIANSON <i>et al.</i> 1992)
pRS315.NDC1	NDC1/CEN/LEU2	(CHIAL <i>et al.</i> 1998)
PSJ906	SPC42- <i>mCHERRY-HIS/LEU2</i>	This Study
PSW863	POM152/2μ/LEU2	(MIAO <i>et al.</i> 2006)
PSW3422	RTN1/CEN/LEU2	(DAWSON <i>et al.</i> 2009)
PSW3673	APQ12/2μ/LEU2	This Study
PSW3674	BBP1/2μ/LEU2	This Study
PSW3675	BRR6/2μ/LEU2	This Study
PSW3676	MPS2/2μ/LEU2	This Study
PSW3592	<i>leu2Δ0::DsRED-HDEL:LEU2</i> integration cassette	This Study

- BEVIS, B. J., A. T. HAMMOND, C. A. REINKE and B. S. GLICK, 2002 De novo formation of transitional ER sites and Golgi structures in *Pichia pastoris*. *Nat Cell Biol* 4: 750-756.
- CHIAL, H. J., M. P. ROUT, T. H. GIDDINGS and M. WINEY, 1998 *Saccharomyces cerevisiae* Ndc1p is a shared component of nuclear pore complexes and spindle pole bodies. *J Cell Biol* 143: 1789-1800.
- CHRISTIANSON, T. W., R. S. SIKORSKI, M. DANTE, J. H. SHERO and P. HIETER, 1992 Multifunctional yeast high-copy-number shuttle vectors. *Gene* 110: 119-122.
- DAWSON, T. R., M. D. LAZARUS, M. W. HETZER and S. R. WENTE, 2009 ER membrane-bending proteins are necessary for de novo nuclear pore formation. *J Cell Biol* 184: 659-675.
- MIAO, M., K. J. RYAN and S. R. WENTE, 2006 The integral membrane protein Pom34p functionally links nucleoporin subcomplexes. *Genetics* 172: 1441-1457.
- SHANER, N. C., R. E. CAMPBELL, P. A. STEINBACH, B. N. GIEPMANS, A. E. PALMER *et al.*, 2004 Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat Biotechnol* 22: 1567-1572.
- SIKORSKI, R. S., and P. HIETER, 1989 A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* 122: 19-27.