


Supplemental Figure 1



Supplemental Figure 1. Adenoviral infection does not induce hepcidin expression. Mice were infected with GFP-expressing adenovirus and analyzed 3 days after infection. Gene expression was examined by qPCR for hepcidin and normalized to β -actin. n = 3 animals per group.

Supplemental Table 1

Primers for qPCR and cloning

qPCR Primers	Sequence (5' to 3')
Human β -actin F	GTTGTCGACGACGAGCG
Human β -actin R	GCACAGAGCCTCGCCTT
Mouse β -actin F	TATTGGCAACGAGCGGTTCC
Mouse β -actin R	GGCATAGAGGTCTTTACGGATGT
Human hepc F	CTCCTTCGCCTCTGGAACAT
Human hepc R	AGTGGCTCTGTTTTCCACA
Mouse EPO F	CATCTGCGACAGTCGAGTTCTG
Mouse EPO R	CACAACCCATCGTGACATTTTC
Mouse C/EBP α F	GAACAGCAACGAGTACCGGGT
Mouse C/EBP α R	GCCATGGCCTTGACCAAGGAG
Mouse Arnt F	CAAGCCATCTTTCCTCACTGATC
Mouse Arnt R	ACACCACCCGTCCAGTCTCA
Mutagenesis Primers	
C/EBP α S193A TOP	ACCCGCACGCGGCTCCCGCGCAC
C/EBP α S193A BOT	GTGCGCGGGAGCCGCGTGCGGGT