

SUPPLEMENT

Regulator of G-Protein Signaling 3 isoform 1 (PDZ-RGS3) Enhances Canonical Wnt Signaling and Promotes Epithelial Mesenchymal Transition

Chong-Shan Shi¹, Ning-Na Huang¹, and John H. Kehrl^{1,2}

¹From the B-Cell Molecular Immunology Section, Laboratory of Immunoregulation, National Institute of Allergy and Infectious Disease, Bethesda, MD 20892, U.S.A

Running Title: *Canonical Wnt signaling and RGS3*

²Correspondence should be addressed to:

John H. Kehrl, B-Cell Molecular Immunology Section, Laboratory of Immunoregulation, National Institute of Allergy and Infectious Disease, 10 Center Drive, MSC 1876, Bethesda, MD 20892, U. S. A. Tel: 301-443-6907. E-mail: jkehrl@niaid.nih.gov

Key words: GSK3 β , epithelial, β -catenin, RGS, signal transduction

Contents:

2 Figures labeled S1 and S2

Figure S1.




Figure S1. Schematic of RGS3 proteins.

Figure S2.




Figure S2. Altered morphology of HEK 293 cells following manipulation of PDZ-RGS3 expression. Shown are images of live HEK 293 cells transfected with the indicated plasmids the previous day. Each image is a superimposed bright field and confocal fluorescent image.