

Model	AIC	C statistics (95% CI)	Hosmer-Lemeshow
<b>CVD</b>			
CKD-EPI			
Model 1	18761.8	0.701(0.638,0.760)	43.2852
Model 2	18477.1	0.711(0.648,0.770)	70.4724
MDRD			
Model 1	18756.9	0.702(0.639,0.761)	44.7680
Model 2	18473.6	0.711(0.649,0.771)	74.4724
<b>CHD</b>			
CKD-EPI			
Model 1	13470.6	0.713(0.640,0.781)	40.2734
Model 2	13221.6	0.721(0.647,0.788)	53.1043
MDRD			
Model 1	13464.7	0.714(0.642,0.782)	37.5166
Model 2	13217.1	0.722(0.649,0.789)	51.5488
<b>Stroke</b>			
CKD-EPI			
Model 1	4600.8	0.717(0.591,0.828)	40.3750
Model 2	4577.8	0.733(0.609,0.842)	53.2918
MDRD			
Model 1	4606.2	0.715(0.589,0.826)	42.6643
Model 2	4581.0	0.733(0.608,0.842)	54.2090
<b>HF</b>			
CKD-EPI			
Model 1	5585.8	0.741(0.631,0.838)	42.4216
Model 2	5463.9	0.758(0.648,0.853)	57.5229
MDRD			
Model 1	5581.7	0.740(0.630,0.837)	46.8696
Model 2	5460.3	0.757(0.647,0.852)	50.1834

**Table S2.** Comparison of Models in Table 2 Using AIC, Discrimination Using C-Statistics, and Calibration Using Hosmer-Lemeshow Statistics.

Abbreviations: AIC = Akaike information criterion; CHD = coronary heart disease; CKD-EPI = Chronic Kidney Disease Epidemiology Collaboration equation; CVD = cardiovascular disease; eGFR = estimated glomerular filtration rate; MDRD = Modification of Diet in Renal Disease equation.