

SUPPLEMENTARY ONLINE DATA



Identification of a haem domain in human soluble adenylate cyclase

Sabine MIDDELHAUFE*†, Martina LEIPELT*, Lonny R. LEVIN†, Jochen BUCK† and Clemens STEEGBORN*§1

*Department of Physiological Chemistry, Ruhr-University Bochum, Bochum, Germany, †Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, U.K., †Department of Pharmacology, Weill Medical College of Cornell University, New York, NY, U.S.A., and §Department of Biochemistry, University of Bayreuth, Bayreuth, Germany

Table S1 Expression constructs for fragments of the human sAC regulatory region fused to \ensuremath{GST}		
Construct (N-terminus \rightarrow C-terminus)	Expression	Solubilit

$\hline \textbf{Construct (N-terminus} \rightarrow \textbf{C-terminus)}$	Expression	Solubility
485–760	_	n.a.
580–760	+	_
870–1091	+	_
897–1081	+	+
897–1130	+	+
965–1130	+	+
1030–1200	+	_
1444–1608	+	_

Received 31 May 2012/18 June 2012; accepted 19 June 2012 Published as Immediate Publication 10 July 2012, doi 10.1042/BSR20120051

 $^{1}\,$ To whom correspondence should be addressed (email Clemens.Steegborn@uni-bayreuth.de).

^{© 2012} The Author(s) This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial Licence (http://creativecommons.org/licenses/ by-nc/2.5/) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.