

TABLE S1. Disc diffusion of randomly collected isolates from Creighton University Medical Center

Drug ^a	<i>E. coli</i> (n = 106)			<i>K. pneumoniae</i> (n = 10)			<i>K. oxytoca</i> (n = 4)		
	S	I	R	S	I	R	S	I	R
AMP	31.9% (23)	5.5% (4)	62.5% (45)	50.0% (2)	0	50.0% (2)	0	0	100.0% (2)
AMC	72.7% (16)	13.6% (3)	13.6% (3)	80.0% (4)	20.0% (1)	0	0	0	0
FEP	98.7% (76)	1.3% (1)	0	0	0	0	0	0	0
CPD	91.4% (64)	2.9% (2)	5.7% (4)	50.0% (4)	0	50.0% (4)	100.0% (1)	0	0
CRO	95.8% (69)	0	4.2% (3)	100.0% (3)	0	0	100.0% (2)	0	0
KAN	80.0% (20)	20.0% (5)	0	100.0% (3)	0	0	0	0	0
CIP	67.2% (45)	3.0% (2)	29.9% (20)	71.4% (5)	0	28.6% (2)	33.3% (1)	0	66.7% (2)
LEX	90.9% (20)	0	9.1% (2)	66.6% (4)	0	33.3% (2)	0	0	0
TZP	95.9% (70)	0	4.1% (3)	60.0% (3)	0	40.0% (2)	0	0	0
SXT	73.1% (49)	0	26.9% (18)	57.1% (4)	0	42.9% (3)	75.0% (3)	0	25.0% (1)
ATM	83.3% (45)	3.7% (2)	13.0% (7)	75.0% (3)	0	25.0% (1)	0	0	0
ETP	100.0% (3)	0	0	0	0	0	0	0	0
GEN	90.9% (40)	0	9.1% (4)	100.0% (3)	0	0	100.0% (2)	0	0
CFZ	43.2% (16)	13.5% (5)	43.2% (16)	0	62.5% (5)	37.5% (3)	75.0% (3)	25.0% (1)	0
SAM	59.5% (25)	11.9% (5)	28.6% (12)	20.0% (1)	40.0% (2)	40.0% (2)	50.0% (1)	50.0% (1)	0
NIT	98.6% (73)	0	1.4% (1)	0	0	0	0	0	0

^aAMP; ampicillin AMC; ampicillin-clavulanate FEP; cefepime CPD; cefpodoxime CRO; ceftriaxone KAN; kanamycin CIP; ciprofloxacin LEX; cephalexin TZP; piperacillin-tazobactam SXT; trimethoprim-sulfamethoxazole ATM; aztreonam ETP; ertapenem GEN; gentamicin; CFZ cephazolin SAM; ampicillin/sulbactam NIT; nitrofurantoin