

SUPPLEMENTAL INFORMATION

Domain Organization of the Polymerizing Mannosyltransferases Involved in Synthesis of the *Escherichia coli* O8 and O9a Lipopolysaccharide O Antigens

Laura K. Greenfield, Michele R. Richards, Evgeny Vinogradov, Warren W. Wakarchuk, Todd L. Lowary, and Chris Whitfield*

*To whom correspondence may be addressed: Chris Whitfield, Department of Molecular and Cellular Biology, Science Complex, University of Guelph, Guelph, Ontario N1G 2W1 Canada. Tel.: 519-824-4120, Ext. 53361; FAX: 519-837-3273; E-mail: cwhitfie@uoguelph.ca.

Table S1: Primer sequences used for the production of recombinant plasmids and site-directed mutants.

Primer	Sequence ^{a,b,c} (5'→3')	Features
WbdCE275AFwd	CGCATCTGCGCTCTGCAGCG TTGGC	Forward primer used to generate the E275A mutation of WbdC ^{O9a} in the creation of pWQ583
WbdCE275ARev	CGTAATGCCAACGCCGCA GAGCGCAG	Reverse primer used to generate the E275A mutation of WbdC ^{O9a} in the creation of pWQ583
WbdCE283AFwd	GGCATTACGTTATTGGCAGG CGCACGC	Forward primer used to generate the E283A mutation of WbdC ^{O9a} in the creation of pWQ584
WbdCE283ARev	CTTGCAAAGCGTGCCTGC CAATAACG	Reverse primer used to generate the E283A mutation of WbdC ^{O9a} in the creation of pWQ584
WbdBE294AFwd	GTTTATCCCTCCTCTATGCG GGATTGGTTTACC	Forward primer used to generate the E294A mutation of WbdB ^{O9a} in the creation of pWQ585
WbdBE294ARev	CAAGAATAGGTAAACCGAA TCCCGCATAGAAGGAGGG	Reverse primer used to generate the E294A mutation of WbdB ^{O9a} in the creation of pWQ585
WbdBE302AFwd	GATTCGGTTTACCTATTCTTG CAGCGATGTCTTGC	Forward primer used to generate the E302A mutation of WbdB ^{O9a} in the creation of pWQ586
WbdBE302ARev	CACACCGCAAGACATCGCTG CAAGAATAGG	Reverse primer used to generate the E302A mutation of WbdB ^{O9a} in the creation of pWQ586
LG33	gatcgaaattcaccatgcacatcatcatcatcatcat catcatcatCATATTGATTGAT GTCCAGGG	Forward primer used for the amplification of <i>wbdA₁₋₁₃₀₅</i> ^{O9a} in the creation of pWQ590; <i>EcoRI</i> restriction site
LG34	gatcaagcttTATTGCTCAGCG CATCGATTCTGC	Reverse primer used for the amplification of <i>wbdA₁₋₁₃₀₅</i> ^{O9a} in the creation of pWQ590; <i>HindIII</i> restriction site
LG35	gatcgaaattcaccatgcacatcatcatcatcat catcatcatcatCAGAAAATCGATG CGCTGAGCG	Forward primer used for the amplification of <i>wbdA₁₂₇₉₋₂₅₂₀</i> ^{O9a} in the creation of pWQ591 and pWQ632; <i>EcoRI</i> restriction site
LG10	gatcaagcttTATTGCTTCGCGG CTGGCGCG	Reverse primer used for the amplification of <i>wbdA₁₂₇₉₋₂₅₂₀</i> ^{O9a} in the creation of pWQ591 and pWQ632; <i>HindIII</i> restriction site
LG101	gatcgaccATGgtACGTATTGTCA TAGATTACAAGGCG	Forward primer used for the amplification of <i>wbdA</i> ^{O8} in the creation of pWQ592, pWQ593 and pWQ595; <i>NcoI</i> restriction site
LG105	gatcgatctagaTTATGCCGTTTC AGATTGTAATC	Reverse primer used for the amplification of <i>wbdA₁₋₁₃₅₀</i> ^{O8} in the

		creation of pWQ592 and pQW593; <i>Xba</i> I restriction site
LG110	<u>gatcgaccATGgtAGCCGAGACGA</u> CAAAAGATGTGG	Forward primer used for the amplification of <i>wbdA</i> ₁₈₄₈₋₃₆₄₂ ⁰⁸ in the creation of pWQ594; <i>Nco</i> I restriction site
LG104	<u>gatcgatctagaTTATTAAAATAT</u> TTAGCTTAGCTAACGCC	Reverse primer used for the amplification of <i>wbdA</i> ⁰⁸ in the creation of pWQ594 and pWQ596; <i>Xba</i> I restriction site
LG102	<u>gatcgatctagaTTAGCCATGAGGT</u> GCGCTTTC	Reverse primer used for the amplification of <i>wbdA</i> ₁₋₂₅₉₅ ⁰⁸ in the creation of pWQ595; <i>Xba</i> I restriction site
LG103	<u>gatcgaccATGgtAGCGCAGGTTT</u> TGCCTGAAGCTTG	Forward primer used for the amplification of <i>wbdA</i> ₁₂₃₄₋₃₆₄₂ ⁰⁸ in the creation of pWQ596; <i>Nco</i> I restriction site
WbdAE317AFwd	GTGTTCCCGTCGCTGCATGC AGGTTTCGGCC	Forward primer used to generate the E317A mutation of WbdA ^{09a} in the creation of pWQ597
WbdAE317ARev	GCGGCAGGCCGAAACCTGC ATGCAGCGACG	Reverse primer used to generate the E317A mutation of WbdA ^{09a} in the creation of pWQ597
WbdAE325AFwd	GCCTGCCGCCGCTGGCAGCG ATCGCTG	Forward primer used to generate the E325A mutation of WbdA ^{09a} in the creation of pWQ598
WbdAE325ARev	CCCGCAGCGCATCGCTGCCA GCGGCGGC	Reverse primer used to generate the E325A mutation of WbdA ^{09a} in the creation of pWQ598
WbdAE750AFwd	CTTTGCCTCGCAGGGAGCAG GCTTTGGC	Forward primer used to generate the E750A mutation of WbdA ^{09a} in the creation of pWQ599 and pLG59
WbdAE750ARev	CGGCAGGCCAAAGCCTGCTC CCTGCGAG	Reverse primer used to generate the E750A mutation of WbdA ^{09a} in the creation of pWQ599 and pLG59
WbdAE758AFwd	GGCCTGCCGTTGATTGCAGC GGCGCAG	Forward primer used to generate the E759A mutation of WbdA ^{09a} in the creation of pWQ630 and pLG60
WbdAE758ARev	GGCAGCTTTCTGCGCCGC TGCAATCAACGGC	Reverse primer used to generate the E759A mutation of WbdA ^{09a} in the creation of pWQ630 and pLG60

^aRestriction sites are underlined.

^bNon-chromosomal sequence is shown in lower-case letters.

^cPoint mutations are bolded.




Figure S1: ¹H-¹³C gHSQC spectrum for the products synthesized by N-WbdA^{O9a} using Acceptor B.