

Additional file 7: Education-related relative inequality in noncommunicable disease prevalence among adults aged 18 or higher living in 41 low- and middle-income countries, World Health Survey 2002-04

		Men						Women					
		Middle-income group			Low-income group			Middle-income group			Low-income group		
		Estimate	95%CI		Estimate	95%CI		Estimate	95%CI		Estimate	95%CI	
Angina	Model 1	1.52	1.19	1.93	2.20	1.70	2.86	1.35	1.10	1.64	1.66	1.35	2.06
	Model 2	1.09	0.83	1.42	1.54	1.14	2.06	1.05	0.87	1.28	1.29	1.01	1.63
Arthritis	Model 1	1.20	0.88	1.63	1.86	1.24	2.80	1.00	0.79	1.27	1.43	1.02	2.00
	Model 1	0.97	0.68	1.39	1.60	1.03	2.50	0.94	0.73	1.20	1.18	0.79	1.75
Asthma	Model 2	2.04	1.47	2.85	2.08	1.52	2.86	1.74	1.12	2.69	1.39	0.98	1.98
	Model 1	1.41	0.99	2.02	1.54	1.02	2.31	1.41	0.95	2.08	1.26	0.87	1.82
Depression	Model 1	1.51	0.96	2.38	1.45	0.91	2.29	1.58	1.07	2.33	2.19	1.68	2.87
	Model 2	1.12	0.65	1.93	1.06	0.63	1.77	1.23	0.81	1.87	1.71	1.22	2.39
Diabetes	Model 1	0.70	0.45	1.08	0.18	0.12	0.29	1.07	0.66	1.72	0.17	0.10	0.31
	Model 2	0.98	0.60	1.61	0.36	0.20	0.64	1.44	0.96	2.17	0.44	0.22	0.88
Co-morbidity	Model 1	1.56	1.10	2.20	2.46	1.78	3.41	1.39	1.04	1.87	1.97	1.44	2.71
	Model 2	1.09	0.74	1.61	1.66	1.12	2.45	1.18	0.90	1.54	1.64	1.13	2.39

* Model 1 is adjusted for country of residence and age

** Model 2 is adjusted for country of residence, age, marital status, urban/rural area and wealth