

Table S1: Sequences of primers used for PCR and sequencing.

Forward (F) and reverse (R) primers are given. Mitochondrial gene loci: CO1 = cytochrome C oxidase 1, CytB = cytochrome B oxidase, 16S = 16S ribosomal RNA. Nuclear gene loci: H3 = histon 3, H4 = histon 4, ARK = arginine kinase, 18S = 18S ribosomal RNA. I = inosine.

| Gene | Primer | F/R | Primer sequence | Reference |
|------|----------|-----|---|------------------------|
| CO1 | Jerry | F | CAA CAT TTA TTT TGA TTT TTT GG | Simon et al. 1994 |
| | Pat | R | TCC AAT GCA CTA ATC TGC CAT ATT A | |
| CytB | CB3 | F | GAG GAG CAA CTG TAA TTA CTA A | Baraclough et al. 1999 |
| | CB4 | R | AAA AGA AA(AG) TAT CAT TCA GGT TGA AT | |
| 16S | M14 | F | CGC CTG TTT AAC AAA AAC AT | Simon et al. 1994 |
| | M223 | R | GGT CCC TTA CGA ATT TGA ATA TAT CCT | |
| H3 | H3aF | F | ATG GCT CGT ACC AAG CAG AC(AG) CGC | Colgan et al. 1998 |
| | H3aR | R | ATA TCC TT(AG) GGC AT(AG) AT(AG) GTG AC | |
| H4 | H4F2s | F | T(CG)C GIG A(CT)A ACA T(CT)C AGG GIA TCA C | Pineau et al. 2004 |
| | H4F2er | R | C(GT)(CT) TTI AGI GC(AG) TAI ACC AC(AG) TCC AT | |
| ARK | AK183F | F | GAT TCT GGA GTC GG(AGCT) AT(CT) TA(CT) GC(AGCT) CC(CT) GA(CT) GC | Wild & Maddison 2008 |
| | AK939R | R | GCC (AGCT)CC (CT)TC (AG)GC (CT)TC (AG)GT GTG (CT)TC | |
| 18S | 18S 5' | F | GAC AAC CTG GTT GAT CCT GCC AGT (I) | Shull et al. 2001 |
| | 18S b5.0 | R | TAA CCG CAA CAA CTT TAA T (I) | |

- Baraclough, T.G., Hogan, J.E., & Vogler, A.P. 1999 Testing whether ecological factors promote cladogenesis in a group of tiger beetles (Coleoptera: Cicindelidae). *Proc. R. Soc. Lond. B.* 266, 1061-1067.
- Colgan, D.J., McLauchlan, A., Wilson, G.D.F., Livingston, S.P., Edgecombe, G.D., Cassis, G. & Gray, M.R. 1998 Histone H3 and U2 snRNA DNA sequences and arthropod molecular evolution. *Aust. J. Zool.* 46, 419-437.
- Pineau, P., Henry, M., Suspène, R., Marchio, A., Dettai, A., Debruyne, R., Petit, T., Lécu, A., Moisson, P., Dejean, A., et al. 2004 A Universal Primer Set for PCR Amplification of Nuclear Histone H4 Genes from All Animal Species. *Mol. Biol. Evol.* 22, 582-588.
- Shull, V.L., Vogler, A.P., Baker, M.D., Maddison, D.R. & Hammond, P.M. 2001 Sequence Alignment of 18S Ribosomal RNA and the Basal Relationships of Aedeopachyinae Beetles: Evidence for Monophyly of Aquatic Families and the Placement of Trachypachidae. *Syst. Biol.* 50, 945-969.
- Simon, C., Frati, F., Beckenbach, A.T., Crespi, B., Liu, H. & Flook, P. 1994 Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Ann. Entomol Soc. Am.* 87, 651-701.
- Wild, A.L. & Maddison, D.R. 2008 Evaluating nuclear protein-coding genes for phylogenetic utility in beetles. *Mol. Phylogenet. Evol.* 48, 877-891.