

Additional file 4. Crude prevalence of risk factors for noncommunicable diseases among adults aged 18 or higher living in 48 low- and middle-income countries, World Health Survey 2002-04

		Current daily smokers			Low-fruit/vegetable consumers <sup>a</sup>			Physically inactive people <sup>b</sup>			Heavy episodic alcohol drinkers <sup>c</sup>		
		Estimate	95%CI		Estimate	95%CI		Estimate	95%CI		Estimate	95%CI	
Men	Middle-income country group	<b>29.0</b>	28.0	29.9	<b>73.7</b>	72.1	75.2	<b>13.0</b>	12.1	14.0	<b>13.6</b>	12.7	14.4
	Low-income country group	<b>29.7</b>	28.3	31.0	<b>74.8</b>	73.0	76.7	<b>6.3</b>	5.5	7.1	<b>3.5</b>	3.1	4.0
Women	Middle-income country group	<b>8.5</b>	7.9	9.0	<b>74.0</b>	72.6	75.5	<b>14.1</b>	13.2	15.0	<b>3.3</b>	2.9	3.6
	Low-income country group	<b>4.7</b>	4.2	5.3	<b>74.7</b>	72.9	76.5	<b>14.1</b>	13.0	15.1	<b>1.0</b>	0.8	1.2

Abbreviations: 95% CI, 95% Confidence Interval

All numbers are in percentage

<sup>a</sup> No data were available for Mexico

<sup>b</sup> No data were available for Morocco and Latvia

<sup>c</sup> Mauritania; and Bosnia-Herzegovina, Comoros, Mauritania and Pakistan were excluded from males and females datasets, respectively