Additional file 6: Interactions between Alpine chamois and domestic ruminants. Numbers refer to the number of game-wardens having reported the corresponding observation, i.e., number of analyzed questionnaires (N) and reported frequency of observations (1: never observed; 2: observed no more than once per year; 3: observed more than once per year).

CHAMOIS with	CATTLE				SHEEP					GOAT			
	N	1	2	3	N	1	2	3	N	1	2	3	
Proximity between species													
Physical contact	40	40	0	0	38	37	1	0	37	33	3	1	
Encounter of less than 50 m	40	10	17	13	38	6	13	19	37	10	12	15	
Encounter of more than 50 m	40	4	9	27	38	3	4	31	37	6	9	22	
Non-simultaneous occupation of the same area	40	7	7	26	38	2	2	34	37	7	6	24	
<b>Duration of encounters</b>													
Encounters of less than 50 m for more than 1 h	40	21	10	9	38	14	12	12	37	23	6	8	
Type of interactions													
Mixing of herds when grazing	40	22	7	11	38	19	7	12	37	23	8	6	
Use of the same natural feeding resources	40	6	11	23	38	4	2	32	37	6	7	24	
Use of the same same salt lick	40	13	11	16	38	9	6	23	37	14	6	17	
Use of the same resting places	40	23	13	4	38	12	8	18	37	15	11	11	
Anthropogenic food sources													
Wildlife supplemental feeding	50	50	0	0	49	49	0	0	48	48	0	0	
Livestock food sources	50	48	2	0	49	45	3	1	48	46	2	0	
Other food sources on private grounds	50	48	1	1	49	47	2	0	48	47	1	0	