

Supplemental Table 1

	Nephrotic (n=49)	Diabetic (n=10)	Nephritic (n=31)	Other (n=9)
Age	55.1 (17.5)*	60.0 (8.4)*	40.1 (13.0)	63.7 (10.6)*
Male (%)	71.4	70.0	61.3	77.8
Creatinine	1.4 (0.9)	2.6 (2.3)	1.5 (1.0)	2.2 (0.7)
GFR	68.3 (33.6)	49.2 (32.0)	64.9 (29.9)	36.8 (23.4)†
Uprotein	8.0 (6.8)	8.6 (7.2)*	3.1 (2.8) †	3.2 (1.9)
%Diabetes	6.1	100.0	3.2	77.8
%Hypertension	71.4	80.0	51.6	11.1
NGAL blot mg/gCr	74.5 (126.3)	315.7 (499.6)	35.3 (45.6)	96.4 (158.8)
NGAL blot (ng/ml)	57.3 (112.0)	164.0 (304.3)	20.0 (21.9)	55.7 (65.1)

* <.01 compared to nephritic † <.01 compared to nephrotic # <.05 compared to nephrotic.

"Other" includes nephroangiosclerosis, ESRD prior to dialysis initiation, light chain nephritis, myeloma, and isolated chronic tubulo-interstitial nephritis.

1057x793mm (72 x 72 DPI)