

Table S1: Summary of characterized *bp* alleles

Allele	Background	Mutagen	Mutation	Reference
<i>bp-1</i>	Ler	EMS	Deletion	Koornneef et al., 1983
<i>bp-2</i>	Ler	FN	Deletion	Douglas et al., 2002
<i>bp-2#</i>	RLD/Ler	T-DNA	Point mutant	Venglat et al., 2002
<i>bp-3</i>	Ler	FN	Deletion	Douglas et al., 2002
<i>bp-4</i>	Ler	EMS	Point mutant	Douglas et al., 2002
<i>bp-5</i>	Col	T-DNA	Deletion	Douglas et al., 2002
<i>bp-6</i>	Ler	EMS	Point mutant	Byrne et al., 2002
<i>bp-7</i>	Ler	EMS	Point mutant	Byrne et al., 2002
<i>bp-8</i>	Ler	EMS	Point mutant	Byrne et al., 2002
<i>bp-9</i>	Col	transposon	Insertion	Kanrar et al., 2006
<i>bp-10</i>	Col	EMS	Point mutant	Ikezak et al., 2010
<i>bp-11</i>	Col	X-rays	Deletion	Redei, unpublished; this work

#the *bp-2* mutant reported by Venglat was isolated concomitantly with the *bp-2* mutant we discovered, hence there are two known *bp-2* mutants.

References:

Byrne ME, Simorowski J, Martienssen RA (2002) ASYMMETRIC LEAVES1 reveals knox gene redundancy. *Development* 129, 1957-1965.

Douglas SJ, Chuck G, Dengler RE, Pelecanda L, Riggs CD (2002) KNAT1 and ERECTA regulate inflorescence architecture in Arabidopsis. *Plant Cell* 14: 547-558.

Ikezaki M, Kojima M, Sakakibara H, Kojima S, Ueno Y, Machida C, Machida Y (2010) Genetic networks regulated by ASYMMETRIC LEAVES1 (AS1) and AS2 in leaf development in Arabidopsis thaliana: KNOX genes control five morphological events. *Plant J* 61, 70–82

Kanrar S, Onguka O, Smith HMS (2006) Arabidopsis inflorescence architecture requires the activities of KNOX-BELL homeodomain heterodimers. *Planta* 224: 1163-1173.

Koornneef M, Vaneden J, Hanhart CJ, Stam P, Braaksma FJ, et al. (1983) Linkage map of *Arabidopsis thaliana*. *J Hered* 74: 265-272.

Venglat SP, Dumonceaux T, Rozwadowski K, Parnell L, Babic V, et al. (2002) The homeobox gene BREVIPEDICELLUS is a key regulator of inflorescence architecture in Arabidopsis. *Proc Natl Acad Sci USA* 99: 4730-4735.