

SUPPLEMENTARY TABLE S1. PLASMA LIPID PROFILE OF APOBEC^{-/-}hApo B-Tg MICE WITH DIFFERENT LDLR ALLELES

	<i>Genotyping (LDLR)</i>	<i>TC (mg/dl)</i>	<i>HDL-C (mg/dl)</i>	<i>TG (mg/dl)</i>	<i>hApoB (mg/dl)</i>	<i>T-Ch/hApoB</i>
Female	WT	173 ± 36	56 ± 3	173 ± 60	86 ± 24	2.1 ± 0.44
	Hetero	268 ± 24	68 ± 3	172 ± 60	114 ± 27	2.4 ± 0.37
	KO	727 ± 107	95 ± 9	391 ± 125	277 ± 46	2.7 ± 0.64
Male	WT	157 ± 25	58 ± 6	222 ± 69	79 ± 22	2.1 ± 0.39
	Hetero	239 ± 53	73 ± 6	233 ± 51	119 ± 57	2.2 ± 0.41
	KO	782 ± 131	100 ± 17	546 ± 207	283 ± 65	2.9 ± 0.66

apobec, gene encoding apolipoprotein B mRNA editing catalytic polypeptide; hApoB-Tg, human apolipoprotein B-transgenic; HDL-C, high-density lipoprotein-cholesterol; hetero, heterozygous; KO, knockout; LDLR, low-density lipoprotein receptor; TC, and T-Ch, total cholesterol; TG, total glycerides; WT, wild type.