

Supplemental Table 1: Pairwise Group Differences in Callosal Volume (Unadjusted for ICV)

	Pairwise Group differences in volume (height) at average age (17.31)									Pairwise Group differences in trajectory (slope)								
	COS vs. SIB			COS vs. NV			SIB vs. NV			COS vs. SIB			COS vs. NV			SIB vs. NV		
	F	df	P	F	df	P	F	df	P	F	df	P	F	df	P	F	df	P
CC Posterior	0.2	1,362	0.6581	0.4	1,362	0.5254	0.85	1,362	0.3565	0.2	3,362	0.8957	0.36	3,362	0.7842	0.12	3,362	0.9502
CC Mid-Posterior	7.41	1,363	0.0068*	3.35	1,363	0.0678	0.6	1,363	0.4398	0.45	3,363	0.7146	2.05	3,363	0.1064	0.32	3,363	0.8141
CC Central	2.91	1,363	0.0892	7.16	1,363	0.0078*	0.56	1,363	0.4547	1.96	3,363	0.1193	0.87	3,363	0.4583	0.99	3,363	0.3954
CC Mid-Anterior	0.18	1,366	0.6704	0.75	1,366	0.693	0.16	1,366	0.693	0.77	2,366	0.4626	0.2	2,366	0.8169	0.3	2,366	0.7412
CC Anterior	0.17	1,366	0.681	0.2	1,366	0.651	0	1,366	0.961	0.1	2,366	0.905	0.22	2,366	0.799	0.27	2,366	0.766
CC Total	0	1,362	0.9991	0	1,362	0.9865	0	1,362	0.9868	0.95	3,362	0.4159	0.05	3,362	0.9863	1.27	3,362	0.2839

COS= Childhood-Onset Schizophrenia probands; NV= healthy comparison subjects; SIB= healthy siblings of Childhood-Onset Schizophrenia probands

Data were covaried for gender at the mean centered age (17.31 years [SD=4.66]).

* None of the results survived correction for multiple comparisons