

Table 1, supplementary material

Night-time and day-time pain responders at the thresholds of $\geq 30\%$ or $\geq 50\%$ for the three treatment cycles of either placebo or FREMS (ITT population).

Night time pain			
	1st cycle	2nd cycle	3rd cycle
Proportion of patients with $\geq 30\%$ VAS night reduction			
n	86	72	57
placebo	10.9 %	35.1 %	27.6 %
FREMS	15.0 %	62.9 %	64.3 %
p-value	0.748	0.019	0.008
Proportion of patients with $\geq 50\%$ VAS night reduction			
n	86	72	57
placebo	4.4 %	18.9 %	24.1 %
FREMS	0.0 %	45.7 %	53.6 %
p-value	0.182	0.015	0.022
Day time pain			
	1st cycle	2nd cycle	3rd cycle
Proportion of patients with $\geq 30\%$ VAS day reduction			
Number of patients	92	67	60
placebo	12.2 %	20.6 %	30.0 %
FREMS	11.6 %	54.6 %	66.7 %
p-value	1.000	0.004	0.004
Proportion of patients with $\geq 50\%$ VAS day reduction			
Number of patients	92	67	60
placebo	6.1 %	14.7 %	23.3 %
FREMS	11.6 %	42.4 %	50.0 %
p-value	0.466	0.015	0.032