

Appendix 3 (as supplied by the authors): Placebo-controlled trials of enoxaparin

Name of author	Date of publication	Duration of therapy (days)	Dosage of enoxaparin	Primary endpoint	Mean age of subjects	Female subjects (%)	Number of subjects
Turpie, et.al(1)	Oct-86	14	30 mg bid	DVT measured with venography	67	52	100
Leclerc, et.al (2)	Jan-92	14	30 mg bid	DVT measured with venography	69	60	131
Kalodiki, et.al (3)	Jun-96	8 - 12	40 mg qd	Composite of DVT measured with venography and PE	67	65	170
Samama, et.al (4)	Jan-97	10 ± 2	40 mg qd	Composite of DVT measured with venography and PE	69	42	93
Fuji, et.al (1)(5)	Jun-08	14	20 mg qd, 20 mg bid, 40 mg qd	Composite of DVT measured with venography and PE	62	88	419
Fuji, et.al (2)(5)	Jun-08	14	20 mg qd, 20 mg bid, 40 mg qd	Composite of DVT measured with venography and PE	70	84	364

Note: qd= quaque die (once daily), bid = bis in die (twice daily), DVT = deep vein thrombosis, PE = pulmonary embolism, VTE = venous thrombo embolic events, RD = risk difference, RR = relative risk, NI = non-inferiority, 1 Source : clinicaltrial.gov

References

1. Turpie AGG, Levine MN, Hirsh J, et al. A randomized controlled trial of a low-molecular-weight heparin (enoxaparin) to prevent deep-vein thrombosis in patients undergoing elective hip surgery. *N Engl J Med* 1986;315:925-9.

2. Leclerc JR, Geerts WH, Desjardins L, et al. Prevention of deep vein thrombosis after major knee surgery--a randomized, double-blind trial comparing a low molecular weight heparin fragment (enoxaparin) to placebo. *Thromb Haemost* 1992;67:417-23.
3. Kalodiki EP, Hoppensteadt DA, Nicolaides AN, et al. Deep venous thrombosis prophylaxis with low molecular weight heparin and elastic compression in patients having total hip replacement. A randomised controlled trial. *Int Angiol* 1996;15:162-8.
4. Samama CM, Clergue F, Barre J, et al. Low molecular weight heparin associated with spinal anaesthesia and gradual compression stockings in total hip replacement surgery. arar study group. *Br J Anaesthes* 1997;78:660-5.
5. Fuji T, Ochi T, Niwa S, Fujita S. Prevention of postoperative venous thromboembolism in japanese patients undergoing total hip or knee arthroplasty: Two randomized, double-blind, placebo-controlled studies with three dosage regimens of enoxaparin. *J Orthopaed Sci* 2008;Sep 1:442-51.