

**SUPPLEMENTAL TABLES**

**Supplemental Table 1. Characteristics of CRIC participants at advanced CKD (defined as eGFR < 20 ml/min/1.73 m<sup>2</sup>)**

<b>Characteristic at advanced CKD</b>	<b>No Echocardiogram (N=175)</b>	<b>Had Echocardiogram (N=638)</b>	<b>P value</b>
<b>Age, year Mean (SD)</b>	59.2 (12.3)	59.0 (11.6)	0.8
<b>Female, %</b>	41.7	49.5	0.07
<b>Black, %</b>	37.1	48.9	0.01
<b>eGFR, ml/min/1.73 m<sup>2</sup> Mean (SD)</b>	17.7 (5.7)	18.3 (4.6)	0.2
<b>Slope of eGFR from entry to advanced CKD, ml/min/1.73 m<sup>2</sup> per year Mean (SD)</b>	-6.9 (5.0)	-6.5 (5.5)	0.4
<b>24H Urine Protein, grams/24 hour Mean (SD)</b>	4.1 (4.5)	2.9 (3.5)	0.003
<b>Body Mass Index, kg/m<sup>2</sup> Median (IQR)</b>	31.2 (26.9, 37.3)	31.9 (26.6, 37.1)	0.6
<b>Hemoglobin, g/dL Mean (SD)</b>	10.9 (1.6)	11.3 (1.6)	0.0004
<b>Systolic blood pressure, mmHg Mean (SD)</b>	140.9 (28.1)	136.9 (25.1)	0.07
<b>Diastolic blood pressure, mmHg Mean (SD)</b>	71.1 (15.0)	70.4 (12.9)	0.6
<b>Ace inhibitor or ARB use, %</b>	56.5	63.6	0.09
<b>Hypertension, %</b>	96.0	97.0	0.5
<b>Diabetes, %</b>	64.0	62.9	0.8
<b>Cardiovascular Disease, %</b>	45.7	45.5	1.0

<b>Characteristic at advanced CKD</b>	<b>No Echocardiogram (N=175)</b>	<b>Had Echocardiogram (N=638)</b>	<b>P value</b>
<b>Peripheral Vascular Disease, %</b>	15.4	11.4	0.2
<b>Congestive Heart Failure, %</b>	18.3	15.4	0.3
<b>Stoke, %</b>	12.6	13.0	0.9
<b>Myocardial Infarction/Prior Revascularization, %</b>	28.6	26.8	0.6
<b>High Cholesterol, %</b>	94.3	95.0	0.7
<b>Current Smoker, %</b>	13.1	13.6	0.9
<b>Alcohol Use, %</b>	46.5	47.2	0.9

**Supplemental Table 2. Characteristics of CRIC participants at ESRD**

<b>Characteristic at ESRD</b>	<b>No Echocardiogram (N=97)</b>	<b>Had echocardiogram (N=218)</b>	<b>P value</b>
<b>Age, year Mean (SD)</b>	59.2 (11.4)	56.0 (12.1)	0.03
<b>Female, %</b>	42.3	46.3	0.5
<b>Black, %</b>	38.1	61.9	<.0001
<b>eGFR, ml/min/1.73 m<sup>2</sup> Mean (SD)</b>	16.7 (3.5)	16.9 (3.5)	0.6
<b>Slope of eGFR from entry to advanced CKD, ml/min/1.73 m<sup>2</sup> per year Mean (SD)</b>	-7.6 (5.7)	-8.6 (5.8)	0.2
<b>24H Urine Protein, grams/24 hour Mean (SD)</b>	3.7 (3.7)	4.0 (4.1)	0.5

<b>Characteristic at ESRD</b>	<b>No Echocardiogram (N=97)</b>	<b>Had echocardiogram (N=218)</b>	<b>P value</b>
<b>Body Mass Index, kg/m<sup>2</sup> Median (IQR)</b>	29.1 (25.9, 34.2)	31.5 (26.6, 36.6)	0.005
<b>Hemoglobin, g/dL Mean (SD)</b>	11.0 (1.6)	11.2 (1.6)	0.3
<b>Systolic blood pressure, mmHg Mean (SD)</b>	145.5 (27.8)	141.0 (26.2)	0.2
<b>Diastolic blood pressure, mmHg Mean (SD)</b>	72.8 (13.3)	72.6 (12.9)	0.9
<b>Ace inhibitor or ARB use, %</b>	58.9	64.4	0.4
<b>Hypertension, %</b>	97.9	98.6	0.6
<b>Diabetes, %</b>	66.0	64.2	0.8
<b>Cardiovascular Disease, %</b>	46.4	46.3	1.0
<b>Peripheral Vascular Disease, %</b>	13.4	8.3	0.2
<b>Congestive Heart Failure, %</b>	17.5	14.2	0.5
<b>Stoke, %</b>	16.5	12.8	0.4
<b>Myocardial Infarction/Prior Revascularization, %</b>	27.8	23.4	0.4
<b>High Cholesterol, %</b>	95.9	94.5	0.6
<b>Current Smoker, %</b>	4.1	5.5	
<b>Alcohol Use, %</b>	18.6	11.5	0.09
<b>Diabetes, %</b>	43.3	50.7	0.2

