Laccase production by *Coriolopsis caperata* RCK2011: Optimization under solid state fermentation by Taguchi DOE methodology

Preeti Nandal¹, Sreenivas Rao Ravella² & Ramesh Chander Kuhad¹*

¹Lignocellulose Biotechnology Laboratory, Department of Microbiology, University of Delhi South Campus, Benito Juarez Road, New Delhi-110021, India

² Institute of Biological Environmental and Rural Sciences (IBERS), Aberystwyth University, Gogerddan, Aberystwyth, Ceredigion, SY23 3EE, UK

* Corresponding Author: Prof. R. C. Kuhad

Lignocellulose Biotechnology Laboratory,

Department of Microbiology,


University of Delhi South Campus,

New Delhi – 110021, India.

(E-mail: kuhad85@gmail.com)


Phone: +91-9871509870 (Mobile)

Fax: +91-11-24115270



Supplementary Figure S1 \mid Optimum performance with the major factors contribution

Variation Reduction Plot (based on Assumed Normal Performance Distributions: Current vs. Improved)



Supplementary Figure S2 | Performance distribution on laccase production by *Coriolopsis* caperata RCK2011 at current and improved conditions under solid state fermentation

Supplementary Figure legends:

Supplementary Figure S1. Optimum performance with the major factors contribution

Supplementary Figure S2. Performance distribution on laccase production by *Coriolopsis*caperata RCK2011 at current and improved conditions under solid state fermentation