

Table S3. Associations of sociodemographic and clinical factors with tuberculosis with an allopatric *Mycobacterium tuberculosis* strain among HIV-infected European patients (n=36).

Variables adjusted for ¹	OR (95% CI)	P value
Male sex	0.85 (0.17-4.17)	0.84
Age groups, years		0.29
16-49	1.0 (ref)	
≥50	0.54 (0.18-1.67)	
Swiss-born	6.64 (0.90-∞)	0.066
Frequent travelling	10.0 (1.69-59.31)	0.011
Immunosuppression ³	1.91 (0.21-∞)	0.60
Contact with foreign-born		0.43 ²
No or unknown contact	1.0 (ref)	
Foreign contact	2.56 (0-99.67)	
Swiss contact	0.50 (0-4.47)	
Degree of immunodeficiency (nadir CD4 cell count cells/μl)		0.36 ²
≥200	1.0 (ref)	
50-199	1.38 (0.21-9.24)	
<50	2.70 (0.33-21.98)	

¹ See Figure S2 for a graphical overview

² P values of linear tests for trend

³ Immunosuppression other than HIV infection (use of anti-TNF blockers, malignancy, organ transplantation, use of steroids or methotrexate)

OR, odds ratio; 95% CI, 95% confidence interval