

Supplemental Table 2. Odds ratios for aerodigestive tract cancer and GAG genotype by site

Site		GAG Genotype					
		GAG-9/9	GAG-7/7	GAG-7/8	GAG-7/9	GAG-8/8	GAG-8/9
Esophagus	<i>No. of Cases:</i>	3	13	2	7	0	0
	<i>OR (95% CI)[†]:</i>	1.00 (ref.)	0.90 (0.23-3.58)	0.46 (0.07-2.97)	0.59 (0.14-2.55)	-	-
Tongue	<i>No. of Cases:</i>	0	16	12	14	2	4
	<i>OR (95% CI)[†]:</i>	-	-	-	-	-	-
Floor of mouth	<i>No. of Cases:</i>	0	4	1	5	0	2
	<i>OR (95% CI)[†]:</i>	-	-	-	-	-	-
Pharynx	<i>No. of Cases:</i>	3	12	8	7	0	1
	<i>OR (95% CI)[†]:</i>	1.00 (ref.)	3.40 (0.41-28.2)	5.67 (0.64-50.3)	2.51 (0.29-21.4)	-	4.12 (0.33-51.3)
Larynx	<i>No. of Cases:</i>	3	19	5	13	0	2
	<i>OR (95% CI)[†]:</i>	1.00 (ref.)	0.51 (0.14-1.87)	0.40 (0.07-2.45)	0.56 (0.15-2.11)	-	0.34 (0.03-3.47)
Other	<i>No. of Cases:</i>	0	16	8	8	3	7
	<i>OR (95% CI)[†]:</i>	-	-	-	-	-	-

Abbreviations: 95% CI, 95% confidence interval; OR, odds ratio.

[†] Odds ratios were adjusted for age, sex, body mass index, alcohol consumption and smoking.