

Table S4. Summary of Edema Severity Over Time – Safety Population

| Visit Category | Placebo (N = 71) n (%) | MSDC-0160 50 mg (N = 71) n (%) | MSDC-0160 100 mg (N = 71) n (%) | MSDC-0160 150 mg (N = 70) n (%) | Pioglitazone 45 mg (N = 71) n (%) | Total (N = 354) n (%) |
|----------------|------------------------------|---|--|--|--|-----------------------------|
| Baseline | N' = 71 | N' = 70 | N' = 71 | N' = 70 | N' = 71 | N' = 353 |
| Absent | 63 (88.7) | 62 (88.6) | 62 (87.3) | 66 (94.3) | 65 (91.5) | 318 (90.1) |
| 1+ | 6 (8.5) | 7 (10.0) | 6 (8.5) | 3 (4.3) | 5 (7.0) | 27 (7.6) |
| 2+ | 2 (2.8) | 1 (1.4) | 1 (1.4) | 0 (0.0) | 0 (0.0) | 4 (1.1) |
| 3+ | 0 (0.0) | 0 (0.0) | 1 (1.4) | 1 (1.4) | 1 (1.4) | 3 (0.8) |
| 4+ | 0 (0.0) | 0 (0.0) | 1 (1.4) | 0 (0.0) | 0 (0.0) | 1 (0.3) |
| Week 4 | N' = 62 | N' = 66 | N' = 66 | N' = 69 | N' = 71 | N' = 334 |
| Absent | 56 (90.3) | 61 (92.4) | 60 (90.9) | 65 (94.2) | 66 (93.0) | 308 (92.2) |
| 1+ | 5 (8.1) | 5 (7.6) | 3 (4.5) | 3 (4.3) | 4 (5.6) | 20 (6.0) |
| 2+ | 1 (1.6) | 0 (0.0) | 2 (3.0) | 0 (0.0) | 0 (0.0) | 3 (0.9) |
| 3+ | 0 (0.0) | 0 (0.0) | 1 (1.5) | 1 (1.4) | 1 (1.4) | 3 (0.9) |
| 4+ | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| Week 12 | N' = 56 | N' = 60 | N' = 62 | N' = 59 | N' = 64 | N' = 301 |
| Absent | 51 (91.1) | 57 (95.0) | 57 (91.9) | 56 (94.9) | 59 (92.2) | 280 (93.0) |
| 1+ | 4 (7.1) | 3 (5.0) | 2 (3.2) | 1 (1.7) | 4 (6.3) | 14 (4.7) |
| 2+ | 1 (1.8) | 0 (0.0) | 2 (3.2) | 1 (1.7) | 0 (0.0) | 4 (1.3) |
| 3+ | 0 (0.0) | 0 (0.0) | 1 (1.6) | 1 (1.7) | 1 (1.6) | 3 (1.0) |
| 4+ | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) | 0 (0.0) |

N' is the number of patients who had an assessment of edema at the specified visit.

Baseline was defined as the measurement at Visit 2. If the Visit 2 measurement was missing, the last valid measurement prior to the first dose of randomized double-blind study medication was used.