

Table S2 Plasmids and strains used in this study.

Plasmid	Description	Resistance*	Reference
pFK78	pML123 expressing GFP	Gm	This work
pML123	Broad host range cloning vector	Gm	Labes et al., 1990
pPL3syl (p63::syl)	Cosmid carrying <i>sylA-E</i> of <i>PsyB301D-R</i>	Tet	Ramel et al. 2009
Strain	Description	Resistance*	Reference
<i>E.coli</i>	Triparental mating pRK600	Cm	R. Dudler
<i>PsyB728a-WT</i>	Isolated from snap beans	Rif	Loper & Lindow, 1999
<i>PsyB728a-ΔsylC</i>	ΔsylC mutant of <i>PsyB728a-WT</i>	RifTet	Groll et al., 2008
<i>PsyB728a-WT-GFP</i>	<i>PsyB728a-WT</i> carrying pFK78	RifGm	This work
<i>PsyB728a-ΔsylC-GFP</i>	<i>PsyB728a-ΔsylC</i> carrying pFK78	RifTetGm	This work
<i>PsyB301D</i>	Isolated from snap beans	Rif	Amrein et al., 2004
<i>PsyB301D ΔsylC</i>	ΔsylC mutant of <i>PsyB301D-R</i>	RifTet	Amrein et al., 2004
<i>PsyB301D ΔsylD</i>	ΔsylD mutant of <i>PsyB301D-R</i>	RifTet	Amrein et al., 2004
<i>PsyB301D ΔsylA-E</i>	ΔsylA-E cluster deletion mutant of <i>PsyB728a</i>	Rif	Ramel et al., 2009
<i>PsyB301D ΔsylA-E(pPL3syl)</i>	<i>PsyB301D ΔsylA-E</i> carrying pPL3syl	RifTet	Ramel et al., 2009
<i>PsyB301D-GFP</i>	<i>PsyB301D</i> carrying pFK78	RifGm	This work
<i>PsyB301D ΔsylC-GFP</i>	<i>PsyB301D ΔsylC</i> carrying pFK78	RifTetGm	This work
<i>PsyB301D ΔsylD-GFP</i>	<i>PsyB301D ΔsylD</i> carrying pFK78	RifTetGm	This work
<i>PsyB301D ΔsylA-E-GFP</i>	<i>PsyB301D ΔsylA-E</i> carrying pFK78	RifGm	This work
<i>PsyB301D Δsyl(pPL3syl)-GFP</i>	<i>PsyB301D ΔsylA-E</i> carrying pPL3syl and pFK78	RifTetGm	This work

* Cm, Chloramphenicol; Gm, Gentamycin; Rif, Rifampicin; Tet, Tetracycline

Amrein, H., Makart, S., Granado, J., Shakya, R., Schneider-Pokorny, J., and Dudler, R. (2004) Functional analysis of genes involved in the synthesis of syringolin A by *Pseudomonas syringae* pv. *syringae* B301D-R. *Mol. Plant-Microbe Interact.* **17**, 90-97.

Groll, M., Schellenberg, B., Bachmann, A. S., Archer, C. R., Huber, R., Powell, T. K., Lindow, S., Kaiser, M., and Dudler, R. (2008) A plant pathogen virulence factor inhibits the eukaryotic proteasome by a novel mechanism. *Nature* **452**, 755-758.

Hwang, M. S. H., Morgan, R. L., Sarkar, S. F., Wang, P. W., and Guttmann, D. S. (2005). Phylogenetic characterization of virulence and resistance phenotypes of *Pseudomonas syringae*. *Appl. Environ. Microbiol.* **71**, 5182-5191.

Labes M., Pühler A., and Simon R. (1990) A new family of RSF1010-derived expression and lac-fusion broad-host-range vectors for gram-negative bacteria. *Gene* **89**, 37-46.

Loper, J. E., and Lindow, S. E. (1987) Lack of evidence for *in situ* fluorescent pigment production by *Pseudomonas syringae* pv. *syringae* on bean leaf surfaces. *Phytopathol.* **77**, 1449-1454.

Ramel, C., Tobler, M., Meyer, M., Bigler, L., Ebert, M. O., Schellenberg, B., and Dudler, R. (2009) Biosynthesis of the proteasome inhibitor syringolin A: the ureido group joining two amino acids originates from bicarbonate. *BMC Biochemistry* **10**:26