

## SUPPLEMENTAL MATERIAL

Dai et al., <http://www.jem.org/cgi/content/full/jem.20122196/DC1>**Table S1.** Characteristics of patients with severe TBI-induced ALI, including age, initial Glasgow coma scale (iGOS), and the time of ALI onset

Patient	Age	iGOS	ALI onset (time after TBI)
	<i>yr</i>		<i>h</i>
1	49	3	6
2	20	4	8
3	40	5	7
4	23	3	6
5	59	3	10
6	33	3	10
7	37	4	9
8	26	3	7
9	52	4	6
10	25	4	8

The mean (SD) for age, iGOS, and ALI onset are 36.4 (14) yr, 3.6 (0.7), and 7.7 (1.6) h, respectively. The range (min-max) for age, iGOS, and ALI onset are 25–59 yr, 3–5, and 6–10 h, respectively.

**Table S2.** Characteristics of patients with lung trauma-induced ALI

Patient	Age	ALI onset (time after lung trauma)
	<i>yr</i>	<i>h</i>
1	52	4
2	51	3
3	44	5
4	50	3
5	32	6
6	33	5
7	41	4
8	33	3

The mean (SD) for age and ALI onset are 42 (8.6) yr and 4.1 (1.1) h, respectively. The range (min-max) for age and ALI onset are 32–52 yr and 3–6 h, respectively.