## **Supporting Information**

## Tobacco mosaic virus rods and spheres as supramolecular high-relaxivity MRI contrast agents

Michael A. Bruckman<sup>1</sup>, Stephen Hern<sup>1</sup>, Kai Jiang<sup>1</sup>, Chris Flask<sup>1,2</sup>, Xin Yu<sup>1,2</sup>, Nicole F. Steinmetz<sup>1,2,3</sup>\*

Department of <sup>1</sup>Biomedical Engineering, <sup>2</sup>Radiology, <sup>3</sup>Materials Science and Engineering, Case Western Reserve University, 10900 Euclid Ave., Cleveland, OH 44106, USA.

\* *Corresponding author:* Prof. Nicole F. Steinmetz, Department of Biomedical Engineering, Radiology, Materials Science and Engineering, Case Western Reserve University School of Medicine, 10900 Euclid Avenue, Cleveland, OH 44106, USA, phone: 216-368-5590, email: nicole.steinmetz@case.edu



Supporting Figure 1 TEM and SEC images of eAlk-TMV, iAlk-TMV and eGd-TMV.



Supporting Figure 2 MALDI-TOF MS of Native-TMV, eAlk-TMV and iAlk-TMV.



**Supporting Figure 3** Before (left) and after (right) images of sucrose gradient density purifications of (from left to right) eAlk-TMV, eGd-TMV, iAlk-TMV, and iGd-TMV.



**Supporting Figure 4** Dynamic light scattering data of iGd-SNPs formed from iGd-TMV used in this paper.



**Supporting Figure 5** NanoDSC analysis of native-TMV and iAlk-TMV showing an increase in stability of iAlk-TMV over native-TMV.



**Supporting Figure 6** SDS-PAGE of denatured TMV samples and subsequent lane analysis to determine the protein concentration of iGd-SNPs. Lane assignments: Native TMV at (1) 1.0

mg/ml, (2) 0.5 mg/ml, (3) 0.25 mg/ml, (4) 0.125 mg/ml, and iGd-SNP at (5) 2x dilution and (6) 4x dilution.



**Supporting Figure 7** Summary comparing relaxivity per volume  $(R_1/V)$  of viral nanoparticle MRI contrast agents. References match paper order.