

Supporting Information for:

Cis–Trans Isomerizations of Proline Residues are Key to Bradykinin Conformations

Nicholas A. Pierson,[†] Liuxi Chen,[‡] David H. Russell,[‡] and David E. Clemmer^{*,†}

[†]*Department of Chemistry, Indiana University, Bloomington, IN 47405*

[‡]*Department of Chemistry, Texas A&M University, College Station, TX 77843*

Supporting Information section includes Figures S1 through S4 as discussed in the main text.




Figure S1. Schematic diagram of the home-built ion mobility – mass spectrometer at Indiana University, described in detail in the Experimental section.




Figure S2. Mass spectrum of acetylated BK. The inset is a 250x blow up of m/z region 366–374, which shows there is no presence of the $[\text{acetyl-BK}+3\text{H}]^{3+}$ ion at m/z 368.2.




Figure S3. IM-IM-MS studies of Pro³→Ala BK (RPAGFSPFR) (red traces), where the IM-MS cross section distribution of [M+3H]³⁺ ions is shown across the bottom, three different IM-IM-MS selections in the middle (vertical dashed line represents where the selection pulse was applied), and IM-IM-MS distributions obtained from collisional activation of the three selections on top. The [BK+3H]³⁺ cross section distribution (dashed traces) is shown in all panels for comparison with the analogue peptide. The IM-IM-MS selection and activation results indicate the middle peak in the IM-MS distribution of Pro³→Ala BK [M+3H]³⁺ is not a stable conformation. Cross section values for the analogue peptide were shifted to compare with BK according to amino acid size parameters in ref. 33.




Figure S4. IM-IM-MS (a) selection of $[\text{Pro}^7 \rightarrow \text{Ala BK} + 3\text{H}]^{3+}$ and (b) activation by application of 60 V ($200 \text{ V} \cdot \text{cm}^{-1}$) in IA2 region. The vertical dashed line represents where across the distribution the selection pulse was applied. Upon activation, the distribution shifts to a higher mobility. Cross section values for the analogue peptide were shifted to compare with BK according to amino acid size parameters in ref. 33.