

Table S4: Bacteria and plasmids

name	relevant characteristics	reference
<i>Escherichia coli</i> ER2566	F ⁻ λ- <i>fhuA2</i> [<i>lon</i>] <i>ompT lacZ</i> ::T7 gene1 <i>gal sulA11</i> Δ(<i>mcrC-mrr</i>)114::IS10 R(<i>mcr-73</i> ::miniTn10-Tet ^S)2 R(<i>zgb-210</i> ::Tn10)(Tet ^S) <i>endA1</i> [<i>dcm</i>]	New England Biolabs
<i>Escherichia coli</i> XL-1 Blue	<i>recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac</i> [F' <i>proAB lacI</i> ^f Δ <i>M15 Tn10 (Tet^r)</i>]	Stratagene
<i>Neisseria meningitidis</i> strain MC58	serogroup B	[1]
<i>Neisseria meningitidis</i> strain 8013	serogroup C	[2]
<i>Neisseria elongata</i> subsp. <i>glycolytica</i> 6171/75 (= ATCC 29315 = NCTC 11050 = DSM 23337)	type strain	[3]
<i>Neisseria mucosa</i> (= ATCC 19696 = NCTC 12978 = DSM 17611)	type strain	[4]
<i>Neisseria mucosa</i> (= ATCC 25996 = NCTC 10774 = DSM 4631)	isolated from human pharyngeal mucosa	[5,6]
<i>Neisseria sicca</i> NRL 30,016 (= ATCC 29259 = DSM 17713)	type strain	[7]
<i>Kingella oralis</i> (ATCC 51147 = DSM 18271)	type strain	[8]
<i>Kingella denitrificans</i> (= ATCC 33394 = NCTC 10995 = DSM 10202)	type strain	[9]
<i>Eikenella corrodens</i> strain 31745	isolated from spinal fluid	[10]
<i>Eikenella corrodens</i> (ATCC 23834 = NCTC 10596 = DSM 8340)	type strain	[11]
p0-DUS	pBluescript II SK+ harbouring <i>pilG</i> ::mTnErm	[12]
pOHA-D4	AT - DUS	this study
pSAF-DV-36	TT - DUS	this study
pOHA-DV-37	AA - DUS	this study
pSAF-DV-38	AT - CCCGTCTGAA	this study
pSAF-DV-39	AT - GACGTCTGAA	this study
pSAF-DV-40	AT - GCAGTCTGAA	this study
pSAF-DV-41	AT - GCCCTCTGAA	this study
pSAF-DV-42	AT - GCCCGCTGAA	this study
pOHA-DV-43	AT - GCCGTGTGAA	this study
pSAF-DV-44	AT - GCCGTCTGGAA	this study
pSAF-DV-45	AT - GCCGTCTCAA	this study
pSAF-DV-46	AT - GCCGTCTGTA	this study
pOHA-DV-47	AT - GCCGTCTGAC	this study
pSAF-DV-48	AG - DUS	this study
pSAF-DV-49	AG - mucDUS	this study
pSAF-DV-50	AG - kingDUS	this study
pSAF-DV-51	AG - eikDUS	this study
pSAF-DV-52	reverse USS	this study
pSAF-DV-53	USS	this study

References

1. Tettelin H, Saunders NJ, Heidelberg J, Jeffries AC, Nelson KE, et al. (2000) Complete genome sequence of *Neisseria meningitidis* serogroup B strain MC58. *Science* 287: 1809-1815.
2. Nassif X, Lowy J, Stenberg P, O'Gaora P, Ganji A, et al. (1993) Antigenic variation of pilin regulates adhesion of *Neisseria meningitidis* to human epithelial cells. *Mol Microbiol* 8: 719-725.
3. Henriksen SD, Holten E (1976) *Neisseria elongata* subsp. *glycolytica* subsp. nov. *International Journal of Systematic Bacteriology* October 1976: 478-481.
4. Veron M, Thibault P, Second L (1959) [*Neisseria mucosa* (*Diplococcus mucosus* Lingelsheim)]. I. Bacteriological description and study of its pathogenicity]. *Ann Inst Pasteur (Paris)* 97: 497-510.
5. Berger U (1971) [*Neisseria mucosa* var. *heidelbergensis*]. *Z Med Mikrobiol Immunol* 156: 154-158.
6. Roberts RJ (1985) Restriction and modification enzymes and their recognition sequences. *Nucleic Acids Res* 13 Suppl: r165-200.
7. Bøvre K (1984) Family VIII. *Neisseriaceae*. In: Krieg NR, Holt JG, editors. *Bergey's manual of systematic bacteriology*. Baltimore: Williams & Wilkins. pp. 288-309.
8. Dewhirst FE, Chen CK, Paster BJ, Zambon JJ (1993) Phylogeny of species in the family Neisseriaceae isolated from human dental plaque and description of *Kingella oralis* sp. nov [corrected]. *Int J Syst Bacteriol* 43: 490-499.
9. Weir S, Lee LW, Marrs CF (1996) Identification of four complete type 4 pilin genes in a single *Kingella denitrificans* genome. *Infect Immun* 64: 4993-4999.
10. Tønjum T, Hagen N, Bøvre K (1985) Identification of *Eikenella corrodens* and *Cardiobacterium hominis* by genetic transformation. *Acta Pathol Microbiol Immunol Scand B* 93: 389-394.
11. Henriksen SD (1969) Corroding bacteria from the respiratory tract. 2. *Bacteroides corrodens*. *Acta Pathol Microbiol Scand* 75: 91-96.
12. Ambur OH, Frye SA, Tønjum T (2007) New functional identity for the DNA uptake sequence in transformation and its presence in transcriptional terminators. *J Bacteriol* 189: 2077-2085.