Supplementary article data

Preoperative waiting time increased the risk of periprosthetic infection in patients with femoral neck fracture

Marianne Westberg¹, Finnur Snorrason², and Frede Frihagen¹

Table 5. Details of clinical data for 17 patients with PJI

Case	Age	Sex	Comorbidity	End result	Time from surgery to death
1	83	М	HT, COPD, iron deficiency anemia. Hospitalized for pneumonia and norovirus gastroenteritis.	Irrigation and debridement	Alive
2	82	F	Recurrent UTI. Treated with AB for UTI post-op.	Irrigation and debridement	Alive
3	87	М	None	Resection arthroplasty Multiple organ failure	1.7 months
4	81	F	Diabetes. Post-op. UTI. Recently operated other hip because of hip fracture	Resection arthroplasty Multiple organ failure	0.8 months
5	75	F	RA. HT	Irrigation and debridement	Alive
6	87	F	Diabetes, coronary heart disease, HT, COPD. Post-op. pneumonia and <i>C. difficile</i> infection	Resection arthroplasty. Severe illness.	4 months
7	77	F	RA, HT, adiposity, liver cirrhosis.	Irrigation and debridement	Alive
8	85	F	RA, coronary heart disease. Post-op. pneumonia and myocardial infarction.	Irrigation and debridement Multiple organ failure	1.5 months
9	71	M	Alcoholism. HT, atrial fibrillation, adiposity.	Antibiotic suppression	Alive
10	77	M	HT, several minor cerbral strokes.	Irrigation and debridement	Alive
11	84	M	Severe Alzheimer's disease, prostate cancer.	Irrigation and debridement	5 months
12	73	F	Diabetes, HT, Bechterew's disease. Hospitalized for pneumonia and pyothorax.	2-stage revision surgery	Alive
13	85	F	Recurrent UTI, asthma, depression, dementia.	Irrigation and debridement	12 months
14	79	F	Hospitalized for incurable cholangiocarcinoma.	Antibiotic suppression	3 months
15	49	M	Alcoholism, liver cirrhosis, obesity, renal failure.	Resection arthroplasty	Alive
16	81	F	HT, Parkinson's disease, depression. Hospitalized for a stable C2 (dens) fracture. Post-op. UTI, <i>C. difficile</i> infection, and severe psychosis.	Antibiotic suppression	6 months
17	79	F	Diabetes, dementia. Recently hospitalized for a myocardial infarction and operated for an elbow fracture. Post-op. pneumonia.	Resection arthroplasty	Alive

HT: hypertension; COPD: chronic obstructive pulmonary disease; UTI: urinary tract infection; AB: antibiotic treatment; RA: rheumatoid arthritis; C. difficile: Clostridium difficile.

¹Department of Orthopaedics, Oslo University Hospital, Oslo; ²Department of Orthopaedics, Drammen Hospital, Drammen, Norway. Correspondence: marianne.westberg@oslo-universitetssykehus.no Submitted 12-07-12. Accepted 12-12-11