

**Table S4. Analysis of bacterial susceptibility to *P. putida* BW11M1.** List of non-*Xanthomonas* bacteria used as indicators.

Bacterial strain	Source or reference*
<b>Gram-negative bacteria</b>	
<b><math>\alpha</math>-Proteobacteria</b>	
<i>Agrobacterium tumefaciens</i> A208	[1]
<i>Azospirillum brasilense</i> Sp7	[2]
<i>Sphingomonas wittichii</i> RW1	[3]
<b><math>\beta</math>-Proteobacteria</b>	
<i>Bordetella avium</i> 197N	[4]
<i>Burkholderia cepacia</i> LMG 1222	BCCM <sup>a</sup>
<i>Burkholderia vietnamiensis</i> LMG 10927	BCCM
<i>Variovorax paradoxus</i> LMG 1797	BCCM
<b><math>\gamma</math>-Proteobacteria</b>	
<i>Aeromonas hydrophila</i> ATCC 7966	ATCC <sup>b</sup>
<i>Citrobacter freundii</i> ATCC 8090	ATCC
<i>Enterobacter aerogenes</i> ATCC 13048	ATCC
<i>Erwinia amylovora</i> CFBP 1430	CFBP <sup>c</sup>
<i>Klebsiella pneumoniae</i> ATCC 13883	ATCC
<i>Proteus vulgaris</i> LMM 2011	LMM <sup>d</sup>
<i>Salmonella enteritidis</i> ATCC 13076	ATCC
<i>Serratia entomophila</i> DSM 12358	DSMZ <sup>e</sup>
<i>Shigella flexneri</i> LMG 10472	BCCM
<i>Yersinia enterocolitica</i> LMG 7899	BCCM
<b>Gram-positive bacteria</b>	
<b>Firmicutes</b>	
<i>Lactobacillus plantarum</i> LMG-P21295	[5]
<i>Lactobacillus rhamnosus</i> GG (LMG 6400)	BCCM
<i>Bacillus megaterium</i> ATCC 13632	ATCC
<i>Bacillus subtilis</i> LMG 7135	BCCM
<i>Staphylococcus aureus</i> ATCC 6358	ATCC
<b>Actinobacteria</b>	
<i>Rhodococcus erythropolis</i> N11	[6]
<i>Mycobacterium smegmatis</i> DSM 43756	DSMZ
<i>Streptomyces coelicolor</i> A3(2)	DSMZ
<i>Streptomyces lividans</i> TK24	[7]

<sup>a</sup> BCCM, Belgian Coordinated Collections of Microorganisms

<sup>b</sup> ATCC, American Type Culture Collection

<sup>c</sup> CFBP: French Collection of Plant-associated Bacteria

<sup>d</sup> LMM: Laboratory of Food Microbiology, KU Leuven

<sup>e</sup> DSMZ: Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures

\* References

1. Wirawan IG, Kang HW, Kojima M (1993) Isolation and characterization of a new chromosomal virulence gene of *Agrobacterium tumefaciens*. J Bacteriol 175: 3208–3212.
2. De Mot R, Vanderleyden J (1989) Application of two-dimensional protein analysis for strain fingerprinting and mutant analysis of *Azospirillum* species. Can J Microbiol 35: 960–967.
3. Miller TR, Delcher AL, Salzberg SL, Saunders E, Detter JC, et al. (2010) Genome sequence of the dioxin-mineralizing bacterium *Sphingomonas wittichii* RW1. J Bacteriol 192: 6101–6102.
4. Sebaihia M, Preston A, Maskell DJ, Kuzniak H, Connell TD, et al. (2006) Comparison of the genome sequence of the poultry pathogen *Bordetella avium* with those of *B. bronchiseptica*, *B. pertussis*, and *B. parapertussis* reveals extensive diversity in surface structures associated with host interaction. J Bacteriol 188: 6002–6015.
5. Ström K, Sjögren J, Broberg A, Schnürer J (2002) *Lactobacillus plantarum* MiLAB 393 produces the antifungal cyclic dipeptides cyclo(L-Phe-L-Pro) and cyclo(L-Phe-trans-4-OH-L-Pro) and 3-phenyllactic acid. Appl Environ Microbiol 68: 4322–4327.
6. Nagy I, Compernolle F, Ghys K, Vanderleyden J, De Mot R (1995) A single cytochrome P-450 system is involved in degradation of the herbicides EPTC (*S*-ethyl dipropylthiocarbamate) and atrazine by *Rhodococcus* sp. strain NI86/21. Appl Environ Microbiol 61: 2056–2060.
7. Simonet JM, Boccard F, Pernodet JL, Gagnat J, Guérineau M (1987) Excision and integration of a self-transmissible replicon of *Streptomyces ambofaciens*. Gene 59: 137–144.