

**Table S2.** Patient and tumor characteristics of *BRCA1* mutation carriers in relation to molecular subtypes

<i>BRCA1</i> tumors	Basal (n = 20)	HER2 (n = 2)	LumA (n = 1)	LumB (n = 9)	Normal (n = 1)
<b>Estrogen receptor</b>					
ER+	2	2	1	9	0
ER-	18	0	0	0	1
<b>Progesterone receptor</b>					
PR+	0	2	1	3	1
PR-	20	0	0	6	0
<b>HER2 status</b>					
HER2+	0	2	0	1	0
HER2-	20	0	1	8	1
<b>Histologic grade</b>					
Grade 1	0	0	0	3	0
Grade 2	4	0	1	1	1
Grade 3	12	2	0	4	0
NA	4	0	0	1	0
<b>Tumor type</b>					
Invasive ductal carcinoma	16	2	1	8	1
Invasive lobular carcinoma	1	0	0	0	0
Mucinous carcinoma	0	0	0	0	0
Medullary carcinoma	2	0	0	0	0
Tubular carcinoma	0	0	0	0	0
Metaplastic carcinoma	0	0	0	0	0
Other	0	0	0	0	0
NA	1	0	0	1	0
<b>Age</b>					
< 50 years	14	2	1	3	1
≥ 50 years	6	0	0	6	0
<b>Menopause status</b>					
Premenopausal	14	2	1	2	1
Perimenopausal	0	0	0	0	0
Postmenopausal	5	0	0	7	0
Other	0	0	0	0	0
NA	0	0	0	0	0