

**Table W1.** Masses of GCTs Used for (A) Immunohistochemistry, Mitoric Figures, TUNEL Assay, Percentage of Necrosis, (B) Tumors from 6-Week-Old Mice Used for Immunofluorescence, and (C) Tumors from 6-Week-Old Mice Used for Western Blot.

(A)		
Age (Weeks)	Treatment Received	Tumor Mass (g, Mean ± SEM)
3	—	0.00466 ± 0.000444
4	0.9% NaCl	0.0515 ± 0.00414
4	5 mg/kg anti-VEGFA	0.0618 ± 0.00936
5	0.9% NaCl	0.178 ± 0.0564
5	5 mg/kg anti-VEGFA	0.0830 ± 0.0163
6	0.9% NaCl	0.723 ± 0.258
6	2.5 mg/kg anti-VEGFA	1.08 ± 0.127
6	5 mg/kg anti-VEGFA	0.307 ± 0.151
6	10 mg/kg anti-VEGFA	0.223 ± 0.0485

  

(B)		
Treatment Received	Tumor Mass (g, Mean ± SEM)	
0.9% NaCl	1.39 ± 0.238	
2.5 mg/kg anti-VEGFA	1.08 ± 0.123	
5 mg/kg anti-VEGFA	0.322 ± 0.0223	
10 mg/kg anti-VEGFA	0.223 ± 0.0485	

  

(C)		
Treatment Received	Tumor Mass (g, Mean ± SEM)	
0.9% NaCl	1.39 ± 0.238	
2.5 mg/kg anti-VEGFA	1.09 ± 0.127	
5 mg/kg anti-VEGFA	0.322 ± 0.0223	
10 mg/kg anti-VEGFA	0.223 ± 0.0485	