

Supplemental Table 1. Differences in individual motor neuropathy scores between fallers/non-fallers and those with and without functional impairment.

Item	Total		Any Fall		No Fall		P
	N	%	N	%	N	%	
37 cramps hands							
High	195	42.0%	29	58.0%	145	39.0%	0.0110
Low	269	58.0%	21	42.0%	227	61.0%	
38 cramps feet							
High	286	61.6%	33	66.0%	225	60.5%	0.4491
Low	178	38.4%	17	34.0%	147	40.0%	
41 Holding a pen							
High	239	51.5%	37	74.0%	180	48.4%	0.0005*
Low	225	58.5%	13	26.0%	192	51.6%	
42 manipulating small objects							
High							0.2058
Low	355	76.7%	42	84.0%	283	76.3%	
	108	23.3%	8	16.0%	88	23.7%	
43 difficulty opening a jar							
High	363	78.2%	43	86.0%	288	77.4%	0.1478
Low	101	21.8%	7	14.0%	84	22.6%	
44 walking bc feet dropped downwards							
High	186	40.3%	30	61.2%	139	37.6%	0.0017*
Low	275	59.7%	19	38.8%	231	62.4%	
45 difficulty walking up stairs because of weakness in your legs							
High							0.0282
Low	321	69.3%	41	82.0%	250	67.4%	
	142	30.7%	9	18.0%	121	32.6%	

Item	Total		Any ADL		No ADL		P
	N	%	N	%	N	%	
37 cramps hands							
High	195	42.0%	61	54.5%	116	37.3%	0.0017*
Low	269	58.0%	51	45.5%	195	62.7%	
38 cramps feet							
High	286	61.6%	78	69.6%	180	57.9%	0.0269
Low	178	38.4%	34	30.4%	131	42.1%	
41 Holding a pen							
High	239	51.5%	83	74.1%	132	42.4%	<0.0001*
Low	225	58.5%	29	25.9%	179	49.2%	
42 manipulating small objects							
High							0.0006*
Low	355	76.7%	97	87.4%	224	72.0%	
	108	23.3%	14	12.6%	87	28.0%	
43 difficulty opening a jar							
High	363	78.2%	103	92.0%	228	73.3%	<0.0001*
Low	101	21.8%	9	8.0%	83	26.7%	
44 walking bc feet dropped downwards							
High	186	40.3%	70	63.1%	101	32.6%	<0.0001*
Low	275	59.7%	41	36.9%	209	67.4%	
45 difficulty walking up stairs because of weakness in the legs							
High	321	69.3%	102	91.9%	191	61.4%	<0.0001*
Low	142	30.7%	9	8.1%	120	38.6%	

\*p<0.0036 (Bonferroni correction for 14 items).