

Table S1: The Rivermead Postconcussion Symptoms Questionnaire.

<b>Symptoms</b>	<b>Controls</b>	<b>Subacute phase</b>		<b>Late phase</b>	
		<b>PCS-</b>	<b>PCS+</b>	<b>PCS-</b>	<b>PCS+</b>
Headaches	0.18 (0.46)	1.40 (1.50)	2.20 (1.30)	0.68 (1.20)	1.80 (1.40)
Feelings of dizziness	0 (0)	0.39 (0.86)	1.80 (1.40)	0.05 (0.23)	0.88 (1.40)
Nausea and/or vomiting	0.03 (0.17)	0.21 (0.81)	1.10 (1.20)	0.13 (0.66)	0.59 (1.00)
Noise sensitivity	0 (0)	0.61 (1.10)	2.10 (1.60)	0.34 (0.85)	1.90 (1.60)
Sleep disturbance	0.26 (0.62)	1.40 (1.70)	2.70 (1.50)	0.47 (1.00)	3.10 (1.20)
Fatigue, tiring more easily	0.41 (0.74)	1.40 (1.60)	3.10 (1.10)	0.34 (0.85)	3.00 (1.10)
Being irritable, easily angered	0.21 (0.73)	0.71 (1.20)	2.40 (1.30)	0.24 (0.79)	1.90 (1.40)
Feeling depressed or tearful	0.06 (0.34)	0.61 (1.20)	1.90 (1.50)	0.05 (0.23)	2.00 (1.40)
Feeling frustrated or impatient	0.21 (0.54)	0.68 (1.20)	2.40 (1.50)	0.11 (0.39)	2.10 (1.40)
Forgetfulness, poor memory	0.44 (0.86)	0.53 (1.00)	2.00 (1.30)	0.32 (0.84)	1.80 (1.40)
Poor concentration	0.56 (0.89)	0.84 (1.20)	2.50 (1.20)	0.53 (1.00)	1.90 (1.30)
Slowness	0.12 (0.41)	0.39 (0.97)	1.80 (1.30)	0.21 (0.62)	1.40 (1.30)
Blurred vision	0.03 (0.17)	0.34 (0.97)	1.50 (1.60)	0.13 (0.58)	1.20 (1.70)
Light sensitivity	0 (0)	0.26 (0.86)	1.60 (1.70)	0.03 (0.16)	1.10 (1.50)
Double vision	0 (0)	0 (0)	0.76 (1.40)	0 (0)	0.18 (0.73)
Restlessness	0.03 (0.17)	0.05 (0.32)	1.20 (1.60)	0 (0)	0.59 (1.10)