

**Supplemental Table 1: Medication class usage by subject group. Medications used in the Psychiatric class were sedative/hypnotic agents (i.e. trazodone and zolpidem).**

Study Group	(1) Allergy/Cold/ENT #	(2) Analgesics #	(3) Antimicrobials #	(4) Asthma/Pulmonary #	(5) Cardiovascular #	(6) Dental/Oral Care #	(7) Dermatologic #	(8) Emergency #	(9) Endocrine/Metabolism #	(10) Gastrointestinal #	(11) Genitourinary #	(12) Hematology/Oncology #	(13) Immunization #	(14) Immunologics #	(15) Neurologic #	(16) Nutrition/Electrolytes #	(17) Obstetrics/Gynecology #	(18) Ophthalmic #	(19) Other #	(20) Psychiatric #	(21) Rheumatologic #
HC	1	5	1	2	12	0	1	0	3	3	6	1	0	1	0	15	0	3	1	0	0
MD	3	5	2	0	8	0	0	0	2	2	3	0	0	0	3	13	0	0	0	1	0
DC	1	3	1	2	20	0	0	0	19	3	3	2	0	0	3	10	0	2	0	1	0
DD	4	5	0	2	19	0	0	0	18	4	3	0	0	0	7	0	2	0	1	0	0
<b>Grand Total</b>	<b>9</b>	<b>18</b>	<b>4</b>	<b>6</b>	<b>59</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>42</b>	<b>12</b>	<b>15</b>	<b>3</b>	<b>0</b>	<b>1</b>	<b>6</b>	<b>45</b>	<b>0</b>	<b>7</b>	<b>1</b>	<b>3</b>	<b>0</b>