

Strain	Males			Females		
	Con/con	Lit/con	Comparison	Con/con	Lit/con	Comparison
<b>C57BL/6J</b>	23.53±0.30	21.09±0.50	<i>t</i> (22)=4.38, <i>p</i> <0.001	17.81±0.26	15.54±0.15	<i>t</i> (22)=7.35, <i>p</i> <0.001
<b>BALB/cJ</b>	26.26±0.29	23.42±0.62	<i>t</i> (22)=4.12, <i>p</i> <0.001	20.79±0.31	18.71±0.42	<i>t</i> (22)=3.94, <i>p</i> <0.001

**Table 1.** Body weights of subjects at the time of FST during the treatment phase of Experiment 1. Data are in grams and are expressed as mean±SEM. n=8-16 per group.

Strain	Males			Females		
	Con/con	Lit/con	Comparison	Con/con	Lit/con	Comparison
<b>C57BL/6J</b>	24.96±0.36	25.15±0.63	$t(22) < 1, NS$	18.33±0.29	17.29±0.19	$t(22) = 3.04, p < 0.01$
<b>BALB/cJ</b>	27.21±0.31	27.30±0.42	$t(22) < 1, NS$	21.28±0.33	20.33±0.44	$t(22) = 1.71, NS$

**Table 2.** Body weights of subjects at the end of the withdrawal phase of Experiment 1. Data are in grams and are expressed as mean±SEM. n=8-16 per group.

<b>Strain</b>	<b>Control</b>	<b>Lithium</b>	<b>Comparison</b>
<b>C57BL/6J</b>	25.45±0.41	21.66±0.45	$t(22)=6.15, p<0.001$
<b>CB6F1</b>	29.94±0.39	24.54±0.50	$t(22)=8.43, p<0.001$

**Table 3.** Body weights of subjects at the end of the Experiment 3. Data are in grams and are expressed as mean±SEM. n=12 per group.