

Table E4. Results from risk reclassification analysis, comparing models with biomarkers of increasing strength to a clinical model where correlation ($\rho = -0.2$) is present between the biomarker and systolic blood pressure.

Odds ratio (OR) of biomarker	Overall NRI (95%CI)	NRI for cases (95%CI)	NRI for controls (95%CI)
1.5	0.019 (0.008, 0.036)	0.020 (0.004, 0.038)	0.000 (-0.002, 0.002)
2.0	0.055 (0.041, 0.077)	0.057 (0.040, 0.079)	-0.002 (-0.011, 0.010)
2.5	0.105 (0.088, 0.130)	0.104 (0.084, 0.129)	0.001 (-0.008, 0.012)
3.0	0.147 (0.129, 0.172)	0.141 (0.123, 0.166)	0.007 (-0.003, 0.017)
3.5	0.183 (0.167, 0.215)	0.169 (0.153, 0.199)	0.014 (0.005, 0.025)
4.0	0.223 (0.206, 0.252)	0.201 (0.185, 0.227)	0.022 (0.011, 0.033)
4.5	0.270 (0.250, 0.298)	0.240 (0.22, 0.266)	0.030 (0.020, 0.042)
5.0	0.288 (0.268, 0.317)	0.252 (0.233, 0.279)	0.036 (0.026, 0.048)
5.5	0.315 (0.294, 0.345)	0.273 (0.252, 0.299)	0.042 (0.031, 0.055)
6.0	0.340 (0.320, 0.371)	0.292 (0.273, 0.316)	0.048 (0.037, 0.063)

[^] The 95% confidence intervals for the NRI using the bootstrap (1000 replications)