

Appendix 1 (as supplied by authors): Cross-sectional association between suicide by a schoolmate and suicidality outcomes in youth, assuming those who reported “I don’t know” regarding exposure to suicide were not exposed, OR (95% CI)

	Unadjusted	Partially Adjusted	Fully Adjusted
<u>Ages 12-13</u>			
Ideation			
Exposure in Past Year	3.39 (2.09-5.50)	3.53 (2.15-5.81)	4.24 (2.37-7.58)
Exposure > Year Ago	3.06 (1.98-4.74)	3.05 (1.97-4.73)	2.62 (1.62-4.25)
No Exposure	Reference	Reference	Reference
Attempt			
Exposure in Past Year	3.14 (1.72-5.71)	3.18 (1.70-5.96)	3.74 (1.78-7.85)
Exposure > Year Ago	3.98 (2.24-7.09)	3.91 (2.20-6.94)	3.34 (1.79-6.23)
No Exposure	Reference	Reference	Reference
<u>Ages 14-15</u>			
Ideation			
Exposure in Past Year	1.74 (1.24-2.46)	1.73 (1.23-2.44)	1.89 (1.28-2.79)
Exposure > Year Ago	1.72 (1.25-2.37)	1.66 (1.21-2.29)	1.55 (1.09-2.22)
No Exposure	Reference	Reference	Reference
Attempt			
Exposure in Past Year	2.40 (1.56-3.70)	2.32 (1.51-3.56)	2.44 (1.49-3.98)
Exposure > Year Ago	2.34 (1.54-3.54)	2.18 (1.44-3.29)	2.12 (1.32-3.41)
No Exposure	Reference	Reference	Reference
<u>Ages 16-17</u>			
Ideation			
Exposure in Past Year	1.88 (1.29-2.74)	1.90 (1.30-2.78)	1.70 (1.07-2.71)
Exposure > Year Ago	1.57 (1.07-2.31)	1.54 (1.04-2.27)	1.80 (1.13-2.86)
No Exposure	Reference	Reference	Reference
Attempt			
Exposure in Past Year	2.27 (1.34-3.84)	2.31 (1.33-4.04)	2.66 (1.35-5.23)
Exposure > Year Ago	1.64 (0.95-2.82)	1.58 (0.91-2.77)	1.60 (0.78-3.27)
No Exposure	Reference	Reference	Reference

*Partially adjusted models account for gender and socioeconomic status. For ages 14-15 and 16-17, fully adjusted models account for gender, socioeconomic status, prior depression/anxiety, prior monthly drinking episodes, and prior drug use. For ages 12-13, fully adjusted models account for gender, SES, and prior depression/anxiety.